
Prairie grasses

identified and described
by vegetative characters

Jan Looman
Research Station, Swift Current, Saskatchewan
Agriculture Canada

Publication 1413 1982



1st printing 1971
Reprinted 1974
Reprinted 1976
Reprinted 1977
Reprinted and enlarged 1982

© Minister of Supply and Services Canada 1982 ✓

Available in Canada through

Authorized Bookstore Agents
and other bookstores

or by mail from

Canadian Government Publishing Centre
Supply and Services Canada
Ottawa, Canada, K1A 0S9

Catalogue No. A 53-1413/1982E Canada: \$9.95
ISBN 0-660-11094-6 Other countries: \$11.95

Price subject to change without notice

CONTENTS

Preface 5

First Section: The Grass Plant 7

Vegetative Parts 7

Other Grasslike Plants 14

Bibliography 15

Glossary 15

Second Section: The Identification and Descriptions of Grasses 19

How to Identify a Grass 19

Vegetative-character Key 19

Descriptions of Species 27

Index to Common Names 242

A rule in centimetres and inches is located on the inside of the back cover.

Preface

Most of the 107 grasses described and illustrated in this book are native species that grow on the rangelands of the Prairie Provinces of Canada. Several have been introduced from other countries and they are grown for hay and pasture on cultivated land. A few are persistent weeds in grain-fields and pastures. Some are abundant throughout the prairie region; others grow only in narrowly defined habitats. Many are valuable forage species in southern British Columbia, Eastern Canada, and Northern United States, as well as in the Prairie Provinces.

It is often necessary to identify a grass before it has flowered, after flowers are no longer present, or, indeed, even after some of the plant has been eaten by animals. Such a need is met by this publication. It provides a key that makes use of vegetative characters only. These characters are present in the roots, stalks, and leaves of grasses.

This book is a revision of Technical Bulletin No. 50, by S. E. Clarke, J. A. Campbell, and W. Shevkenek, published in 1944 and reprinted as Publication No. 762 in 1950. It was entitled *The Identification of Certain Native and Naturalized Grasses by their Vegetative Characters*. Contributions are gratefully acknowledged from the original publication and from *The Identification of Certain Native and Naturalized Hay and Pasture Grasses by their Vegetative Characters*, by F. S. Nowosad, D. E. Newton Swales, and W. G. Dore, Macdonald College Technical Bulletin No. 16, 1946.

In the 1971 edition, by Keith F. Best, Jan Looman, and J. Baden Campbell (and under the present title), diagrammatic drawings were added, the key was simplified, and all the descriptions were rewritten. In the present reprinting three species have been added.

Since the last reprinting in 1977 K. F. Best and J. B. Campbell have retired, and Mr. Campbell is now deceased. The continuing need for this publication is a tribute to their work.

THE GRASS PLANT

It is strange that grass, the most common plant in most places in the world, is intimately known to rather few people. The structure of a grass is, however, just as easy to understand as that of any other plant. A typical grass is shown in Figure 1.

The *roots* of grasses, like those of other land plants, serve to anchor the plant in the soil and to conduct water and nutrients in solution from the soil to the plant parts. The main body of a grass comprises the *culm*, or stalk, and the *leaves*. In most grasses the culm is hollow and is composed of several tubes closed at the joints by solid tissue. The joints are *nodes*; the portions between them are *internodes*. One or more buds, which are undeveloped leaf or flower branches, may form at every node.

The leaves are arranged in two ranks, that is, alternately on opposite "sides" of the culm. The base of each leaf is at a node. The expanded upper part of the leaf is the *blade*; the lower part, which is usually split on the "side" opposite the blade, surrounds the culm, and is called the *sheath*.

At the junction of the sheath and blade, and facing the culm, there is almost always a small appendage termed the *ligule*. At the same place, but on the outer or lower "side", there is a band called the *collar*. Usually, on either side of the split in the sheath the collar bears projections termed *auricles*.

An important feature in the identification of grasses is the *emerging leaf*, that is, the new growth as it emerges from the sheath. The emerging leaf is described on page 13.

A description of the flowers is not needed here, because in this book grasses are identified by vegetative characters alone.

A fuller account of essential vegetative characters follows.

VEGETATIVE PARTS

Roots

In grasses there are mainly three types of roots (Figure 2). A fibrous root branches in all downward directions from the crown, as in crested wheat grass (*Agropyron cristatum*) or rough fescue (*Festuca campestris*). Rhizomatous roots, or rootstalks, are actually underground stems from which stalks and roots develop at intervals, as in western wheat grass (*Agropyron smithii*), smooth brome (*Bromus inermis*), and some other grasses. Stolons are horizontal stalks that root at the nodes. The roots of creeping bent (*Agrostis stolonifera* var. *major*) are stoloniferous.




Figure 1 A grass plant, showing the vegetative parts.

Blade

The blade (Figure 3) is usually long and narrow. Though ordinarily rather flat, it may be V-shaped, folded, or rounded. When it is so tightly folded that it seems to be solid and cylindrical, it is termed bristle-like. A rounded leaf may be U-shaped, cylindrical with one margin overlapping the other (convolute), or cylindrical with margins turned inward toward the midrib (involute). A blade may be constricted at the base, or sometimes twisted.




Figure 2 Types of grass roots.

Its tip may be sharp-pointed or boat-shaped, and the general shape may be tapered or mostly with parallel sides (Figure 4). The surface may be smooth, rough, or hairy, and the margins smooth, without hairs (glabrous), or with a fringe of hairs (ciliate). Veins may be so prominent on the upper surface that they form ridges. On the lower surface the midvein or midrib may be prominent enough to form a keel.




Figure 3 Shapes of blades, in cross section.

Figure 4 Shapes of blades and tips.






Figure 5 Split and unsplit sheaths, viewed directly and in cross section.

Figure 6 Ligules, showing (upper row) types (and, right, a leaf without a ligule), (middle row) shapes, and (lower row) margins.




Sheath

The sheath may be round or compressed. Occasionally it is keeled at the midrib. It is usually of a paler shade of green than the blade, and is often tinged with pink or purple at its base. These tints are not consistent enough to be of much value in identification, except in a few grasses such as meadow foxtail (*Alopecurus pratensis*). A sheath may be split completely, split with one margin overlapping the other, or without a split (Figure 5). A ruptured sheath must be distinguished from one that is naturally split.

Ligule

The ligule may be a membrane or a fringe of hairs. In some species the ligule is not present. In texture and color it varies from thin and white, as in Canby blue grass (*Poa canbyi*), to thick and opaque, as in fringed brome (*Bromus ciliatus*). The length of a ligule may vary within a species, but its shape and color and the character of its margins are usually quite constant. The outer surface of the ligule is usually smooth, but it may be pubescent, as in reed canary grass (*Phalaris arundinacea*) and some species of *Agrostis*. The types, shapes, and margins of ligules are shown in Figure 6. Large ligules and those in old or dry material may be so frayed and broken that they are useless for identification purposes.


Figure 7 Types of collars.



Collar

The band that forms the collar (Figure 7) may be vertically broad or narrow, continuous, or divided by a midrib. If it extends diagonally it is called oblique. The collar is usually smooth, but it may be hairy (pubescent) over the whole surface, as in quack grass (*Agropyron repens*), or it may have minute hairs (cilia) on the inner and lowermost portion of the margin. It is usually pale green or yellowish green, but it may be temporarily tinged with red, as in yellow foxtail (*Setaria glauca*).

Figure 8 Shapes of auricles and (right) a collar without auricles.



Auricles

Auricles (Figure 8) are appendages extending from the collar. They may be horizontally clawed, fully or slightly rounded, or rudimentary. Sometimes they are not present. When old or dry specimens are being examined care must be taken to avoid breaking off the auricles.

Emerging Leaf

A developing leaf is closely surrounded by the sheath of the previously developed leaf (Figure 9). In the developing state, a blade may be conduplicate, that is, folded at the midvein with the margins meeting, or it may be rolled lengthwise with either one margin overlapping the other (convolute) or both margins turned inward toward the midrib (involute). Conduplicate leaves usually form a laterally compressed shoot, rolled leaves usually a cylindrical one. There are exceptions. In yellow foxtail (*Setaria glauca*) and barnyard grass (*Echinochloa crusgalli*) the leaves are rolled in flattened shoots, whereas in poverty oat grass (*Danthonia spicata*), June grass (*Koeleria gracilis*), and certain fescues, the folded leaves are in round shoots. In plains reed grass (*Calamagrostis montanensis*) and Nuttall's alkali grass (*Puccinellia nuttalliana*) the leaves are involute in somewhat rounded shoots.

The shape of the emerging leaves can best be seen when the stalk is cut just below the ligule and examined with a hand lens.




Figure 9 Shapes of emerging leaves, with, at right of each leaf, a cross-section view at the juncture of leaf blade and sheath.

OTHER GRASSLIKE PLANTS

Some common plants, mainly the sedges and rushes, are in certain ways similar to grasses. Table 1 presents comparative information.

TABLE 1 A COMPARISON OF PLANT PARTS IN THE GRASSES, SEDGES, AND RUSHES

Character	Gramineae (grasses)	Cyperaceae (sedges)	Juncaceae (rushes)
Culm	Usually hollow; cylindrical or flattened Nodes conspicuous	Filled with pith, rarely hollow, usually three-sided Nodes indistinct	Filled with spongelike pith, cylindrical Nodes indistinct
Leaf arrangement	Two-ranked	Three-ranked	Three-ranked
Leaf blade	Usually flat; often folded, involute, or bristle-like; glabrous or pubescent	Flat, folded, or bristle-like; rarely pubescent	Channeled or round, usually glabrous
Leaf margins	Smooth, scabrous, or ciliate	Usually scabrous	Smooth
Leaf sheath	Usually split, occasionally closed	Usually closed	Open or closed
Ligule	Usually present	Absent or weakly developed	Absent or weakly developed
Collar	Distinct	Indistinct	Indistinct
Auricles	Present or absent	Absent	Absent

BIBLIOGRAPHY

- Chase, Agnes. 1959. First book of grasses. Smithsonian Institution, Washington, D.C. 127 p.
- Hitchcock, A. S. 1951. Manual of the grasses of the United States. Second edition, revised by Agnes Chase. United States Department of Agriculture Miscellaneous Publication No. 200. United States Government Printing Office, Washington, D.C. 1051 p.
- Hubbard, W. A. 1955. The grasses of British Columbia. British Columbia Provincial Museum, Department of Education Handbook No. 9, Victoria, B.C. 204 p.
- Looman, J. 111 Range and forage plants of the Canadian prairies. Agric. Can. Publ. In press.
- Looman, J.; Best, K. F. Budd's flora of the Canadian Prairie Provinces. Agric. Can. Publ. 1662; 1979; 863 p.
- Moss, E. H. 1959. Flora of Alberta. University of Toronto Press, Toronto. 546 p.
- Nowosad, F. S., D. E. Newton Swales, and W. G. Dore. 1946. The identification of certain native and naturalized hay and pasture grasses by their vegetative characteristics. Macdonald College Technical Bulletin No. 16. Macdonald College, Que. 78 p.
- Scoggan, H. J. 1957. Flora of Manitoba. National Museum of Canada Bulletin No. 140, Biological Series No. 47. Canada Department of Northern Affairs and National Resources, Ottawa. 619 p.
- United States Forest Service. 1937. Range plant handbook. Prepared by Forest Service, United States Department of Agriculture. Government Printing Office, Washington, D.C. 842 p.

GLOSSARY

Definitions are limited to the sense of the word as it applies to vegetative parts of grasses in this publication.

- acuminate *of a blade, ligule, stolon, or rhizome: gradually tapering to a sharp point*
- acute *of a blade or ligule: abruptly sharp-pointed*
- annual *of a plant: (noun) a plant that completes its life cycle in one growing season (adjective) completing the life cycle in one growing season*
- ascending *of a plant part: sloping upward*
- auricle *an appendage of a collar*
- blade *the portion of the leaf above the sheath*
- bulbous *of a stalk: having a bulb-like base*
- chartaceous *of a ligule: having the texture of writing paper*
- ciliate *of a blade, collar, or ligule: fringed with fine hairs*
- ciliolate *of a ligule or blade margin: fringed with minute hairs*
- clasping *of an auricle: having the appendage extending directly into the collar and partly or completely surrounding the blade*
- collar *the band on the outer or lower "side" of the leaf at the junction of the sheath and the blade*

compressed	<i>of a sheath:</i> flattened laterally
conduplicate	<i>of a leaf:</i> folded together lengthwise in the emerging leaf
constricted	<i>of a collar:</i> narrowed at the midpoint
continuous	<i>of a collar:</i> extending from one margin of the sheath to the other
convolute	<i>of a leaf:</i> rolled lengthwise in the emerging leaf, with one margin overlapping the other
coriaceous	<i>of a leaf:</i> leathery in texture
corrugated	<i>of a ligule:</i> having alternating ridges and grooves
crown	<i>of a plant:</i> the region from which both stalks and roots grow
decumbent	<i>of a stalk:</i> curved upward from a horizontal or slightly inclined base
divided	<i>of a collar:</i> separated at the midrib
emarginate	<i>of a ligule:</i> notched at the tip
entire	<i>of a blade or ligule:</i> having the margins continuous and not in any way divided
fascicle	<i>of leaves:</i> a compact cluster
fibrillose	<i>of a root:</i> having fine fibers
filiform	<i>of a blade:</i> very slender; threadlike
flaccid	<i>of a blade:</i> lax and weak; lacking rigidity
flexuous	<i>of a blade:</i> having alternate opposite curvatures
glabrous	<i>of a blade, collar, or sheath:</i> without hairs
glaucous	<i>of a blade:</i> covered with a waxy layer, which gives the plant part a bluish green color
hirsute	<i>of a blade or sheath:</i> having coarse, straight, rather stiff hairs
hispid	<i>of a blade or sheath:</i> having stiff hairs or bristles
hyaline	<i>of a sheath margin:</i> thin and translucent or transparent
hybrid	<i>of a plant:</i> a cross between two species of plants
involute	<i>of a blade or an emerging leaf:</i> having margins rolled over the upper surface toward the midrib
keel	a ridge on the back of a sheath or blade usually along the midrib
lacerate	<i>of a ligule:</i> having margins deeply and irregularly cut
ligule	a thin appendage projecting from the inner surface of a leaf at the junction of the sheath and blade
linear	<i>of a blade:</i> long and narrow with parallel sides
membranous	<i>of a ligule:</i> thin, rather soft, and more or less translucent and pliable
midrib	<i>of a blade:</i> the central vein
muriform	<i>of cells:</i> arranged like rows of bricks

oblique	<i>of a collar:</i> slanting and of varying width
obtuse	<i>of a ligule:</i> blunt or rounded at the tip
papillose	<i>of a blade or sheath:</i> having minute nipple-shaped projections
perennial	<i>of a plant:</i> (noun) a plant that lives for more than two years (adjective) living more than two years
pilose	<i>of a plant part:</i> having soft, fairly long, straight hairs
procumbent	<i>of a stalk:</i> partly prostrate
prostrate	<i>of a stalk:</i> trailing along the ground
puberulent	<i>of a sheath:</i> having fine minute hairs
pubescent	<i>of a plant part:</i> having soft fine hairs
retorse	<i>of hairs:</i> bent downward or backward
rhizome	an underground stalk from which, at intervals, shoots arise above and roots descend below; also called root-stalk
ridged	<i>of a blade:</i> having raised veins
rootstalk	a rhizome
rudimentary	<i>of a plant part:</i> imperfectly developed
scaberulous	<i>of a blade:</i> having minute protuberances that are rough to the touch
scabrous	<i>of a blade:</i> having small protuberances that are rough to the touch
scarious	<i>of a sheath or blade:</i> having rather thin, translucent, not green, dry margins
serrate	<i>of a ligule or blade margin:</i> having sharp teeth
serrulate	<i>of a ligule:</i> having fine teeth
sheath	the lower part of the leaf that surrounds the stem or shoot
stolon	an aboveground creeping stem that roots at the nodes
stoloniferous	<i>of a plant:</i> bearing stolons
stoma	(plural: stomata) a minute opening in the surface of a leaf
striate	<i>of a sheath:</i> having very narrow longitudinal lines of darker or lighter color than the adjacent tissue
truncate	<i>of a ligule:</i> having the tip seemingly cut off at a right angle to the midrib
tufted	<i>of plants:</i> having shoots in a loose, compact, or dense cluster arising from a crown
undulate	<i>of a ligule:</i> having a wavy summit
vein	<i>of a blade or sheath:</i> one of the parallel structures that serve to support the leaf
villose	<i>of a plant part:</i> having long, soft hairs

THE IDENTIFICATION AND DESCRIPTION OF GRASSES

HOW TO IDENTIFY A GRASS

The following key contains a series of two or more contrasting statements. Compare the characteristics of the grass you want to identify with the distinguishing feature(s) mentioned in the statements. If the character(s) are present in your grass, and the statement does not lead directly to a species name, notice the number at the extreme *right* of the statement. Go to this number at the *left* of the page and read the adjoining statement(s). Follow this procedure until you read a statement that is followed by a botanical name. This is the name of the grass you have been trying to identify.

In using the key you may need to refer to the glossary, which immediately precedes this section. Note also that for the purpose of this key and in the descriptions of species the word *blades* implies and refers only to mature blades near the collar. To distinguish some characters you may need to use a ten-power hand lens. Individual characters may show some variation. Such variability, as well as changes brought about by shrinkage or distortion in dried or old material, should be taken into account when you use the key.

VEGETATIVE-CHARACTER KEY

1 Auricles present	2
Auricles absent or rudimentary	14
2 Plants with rhizomes	3
Plants without rhizomes	7
3 Rhizomes long, white; blades stiff, glaucous; collar and auricles usually purplish	<i>Agropyron smithii</i>
Rhizomes long, yellowish white; blades not stiff, with lines of stomata on lower surface ..	<i>Agropyron repens</i>
Rhizomes and blades not as above	4
4 Blades less than 10 mm wide	5
Blades more than 10 mm wide	6
5 Rhizomes numerous; blades to 6 mm wide, flat or involute, stiff, dull green	<i>Agropyron dasystachyum</i>
Rhizomes few; blades to 8 mm wide, flat, lax, glossy green beneath; ligule entire	<i>Festuca elatior</i>
Rhizomes few; blades to 10 mm wide, pale green; ligule serrate	<i>Agropyron intermedium</i>

6 Rhizomes short; often absent in old plants; ligule truncate; collar broad, continuous	<i>Elymus canadensis</i>
Rhizomes slender and scaly; ligule truncate; collar narrow, indistinct, divided	<i>Elymus innovatus</i>
Rhizomes short, thick; ligule obtuse, lacerate; collar narrow	<i>Elymus cinereus</i>
7 Blades more than 10 mm wide	8
Blades less than 10 mm wide	9
8 Sheath with hyaline margins; collar broad; ligule very short, about 0.5 mm; blades green	..	<i>Elymus virginicus</i>
Sheath with scarious margins; collar broad; ligule to 2 mm; blades green	<i>Elymus canadensis</i>
Sheath with scarious margins; collar narrow; ligule to 2 mm; blades glaucous	<i>Elymus glaucus</i>
9 Plants annual; blades acuminate, twisted, bluish green	<i>Lolium persicum</i>
Plants perennial; blades not twisted	10
10 Blades soft pubescent on upper surface	<i>Agropyron cristatum</i>
Blades not pubescent	11
11 Sheath usually compressed, reddish at base; blades acuminate, glossy beneath	<i>Lolium perenne</i>
Sheath not compressed, margins hyaline	12
Sheath not compressed, margins scarious	13
12 Collar broad, continuous; blades prominently veined; no lines of stomata on lower surface; ligule truncate	<i>Elymus junceus</i>
Collar broad, continuous, constricted at midrib; blades strongly veined, firm, rigid, with lines of stomata on lower surface; ligule acute	<i>Elymus angustus</i>
Collar distinct, continuous; blades firm, midrib prominent near collar; ligule serrate, very short	<i>Agropyron elongatum</i>
Collar indistinct; blades not prominently veined; ligule truncate	<i>Agrohordeum macounii</i>
13 Plants densely tufted; collar continuous; midrib prominent on lower surface of blade; sheath glabrous	<i>Agropyron spicatum</i>
Plants loosely tufted; collar divided; midrib not prominent on lower surface of blade; sheath pubescent	<i>Agropyron subsecundum</i>
14 Plants with rhizomes	15
Plants without rhizomes; roots fibrous	45
15 Tips of blades boat-shaped	16
Tips of blades acute or acuminate	19
16 Ligule to 5 mm long, white, acuminate; sheath with crossveins	17
Ligule to 3 mm long, greenish, truncate to acute; sheath without crossveins	18
17 Blades narrow, to 6 mm wide; sheath tinged with purple; blades not conspicuously crossveined	<i>Glyceria striata</i>

Blades broad, to 15 mm wide; sheath pale green; blades conspicuously crossveined	<i>Glyceria grandis</i>
18 Sheath strongly flattened, sharply keeled; collar divided; blades bluish green	<i>Poa compressa</i>
Sheath not strongly flattened or keeled; collar continuous; blades glossy dark green	<i>Poa pratensis</i>
Sheath not strongly flattened or keeled; collar usually continuous; blades bluish green	...	<i>Poa glaucifolia</i>
19 Blades more than 10 mm wide	20
Blades less than 10 mm wide	25
20 Sheath closed to near the top	21
Sheath split	22
21 Midrib of blade extending partway down sheath; blade and sheath often pubescent; sheath purplish at base	<i>Bromus pumpeianus</i>
Midrib of blade not extending down sheath; blade and sheath glabrous; sheath not purplish at base	<i>Bromus inermis</i>
22 Plants of dry habitats	23
Plants of wet habitats	24
23 Sheath margins hyaline; sheath usually somewhat pubescent above, light green	<i>Calamovilfa longifolia</i>
Sheath margins scarious and villose; sheath glabrous, strongly veined	<i>Panicum virgatum</i>
24 Ligule membranous, to 5 mm long, white, acute or obtuse; blades to 15 mm wide	<i>Phalaris arundinacea</i>
Ligule a fringe of hairs; blades to 15 mm wide, distinctly veined, smooth and shiny below	<i>Spartina pectinata</i>
Ligule a fringe of hairs; blades to 30 mm wide, not distinctly veined, dull below	...	<i>Phragmites communis</i>
25 Rhizomes very short or absent	26
Rhizomes elongated	33
26 Emerging leaves folded	27
Emerging leaves rolled	28
27 Blade strongly veined; ligule very short; collar indistinct; plants loosely tufted	<i>Festuca rubra</i>
Blade strongly veined; ligule very short; collar indistinct; plants densely tufted	<i>Festuca hallii</i>
Blade not strongly veined; ligule to 2 mm long; collar broad, continuous	<i>Andropogon scoparius</i>
28 Ligule a fringe of hairs	29
Ligule membranous	30
29 Blades to 7 mm wide, twisted; sheath usually pubescent	<i>Bouteloua curtipendula</i>
Blades to 3 mm wide, usually strongly curled; sheath glabrous or somewhat pilose	<i>Bouteloua gracilis</i>
30 Sheath compressed or flattened, pubescent or scabrous	31
Sheath round, glabrous	32

31 Plants densely tufted, blades to 3 mm wide; ligule very short, ciliate	<i>Sporobolus heterolepis</i>	
Plants not densely tufted; blades to 10 mm wide; ligule to 3 mm long, ciliate or lacerate	<i>Andropogon gerardii</i>	
32 Blades basal, glossy green, usually convolute..	<i>Oryzopsis asperifolia</i>	
Blades erect or drooping, dull green, flat..	<i>Agropyron trachycaulum</i>	
33 Emerging leaves folded	34	
Emerging leaves rolled	37	
34 Plants aquatic; blades soft and flat	<i>Catabrosa aquatica</i>	
Plants not aquatic; blades not soft	35	
35 Ligule a fringe of hairs; sheath with scarios margins; plants sod-forming	<i>Distichlis stricta</i>	
Ligule membranous; sheath with hyaline margins; plants tufted	36	
36 Blades to 6 mm wide; rather lax, to 15 cm long; plants loosely tufted	<i>Muhlenbergia racemosa</i>	
Blades to 2 mm wide, rather stiff, rarely more than 5 cm long; plants loosely tufted	<i>Muhlenbergia asperifolia</i>	
Blades to 2 mm wide; rather stiff, to 10 cm long; plants densely tufted	<i>Muhlenbergia richardsonii</i>	
37 Ligule a fringe of hairs	<i>Spartina gracilis</i>	
Ligule membranous	38	
38 Sheath closed in early stages, splitting in middle at maturity, purplish at base ...	<i>Schizachne purpurascens</i>	
Sheath split	39	
39 Sheath with scarios margins	40	
Sheath with hyaline margins	42	
40 Plants with distinctive odor when crushed	<i>Hierochloe odorata</i>	
Plants without odor when crushed	41	
41 Blades rather lax, to 10 mm wide; collar glabrous	<i>Calamagrostis canadensis</i>	
Blades lax, to 4 mm wide; collar pubescent	<i>Calamagrostis rubescens</i>	
Blades stiff, to 4 mm wide; collar glabrous	<i>Calamagrostis inexpansa</i>	
42 Blades narrow, usually less than 5 mm wide	43	
Blades usually more than 5 mm wide	44	
43 Plants of dry prairie; ligule acute, to 7 mm long; blades bluish green	<i>Calamagrostis montanensis</i>	
Plants of marshland; ligule truncate, to 4 mm long; blades green	<i>Calamagrostis neglecta</i>	
44 Plants tufted; blades short, usually less than 15 cm long; woodlands and moist prairie	<i>Phleum alpinum</i>	
Plants not tufted; blades to 50 cm long; moist prairie	<i>Sorghastrum nutans</i>	
Plants not tufted; blades to 35 cm long; wetland and slough margins	<i>Scolochloa festucacea</i>	
45 Ligule a fringe of hairs or absent	46	
Ligule membranous	54	

46 Blades 10 mm or more wide; annuals	47	
Blades less than 10 mm wide	49	
47 Ligule absent; sheath compressed, keeled; shoots prostrate or ascending	<i>Echinochloa crusgalli</i>	
Ligule present; sheath compressed or round; shoots erect	48	
48 Sheath compressed, keeled, glabrous; blades pubescent at base	<i>Setaria glauca</i>	
Sheath compressed, not keeled, pubescent; blades not pubescent at base	<i>Setaria viridis</i>	
Sheath round, conspicuously pubescent; blades pubescent on both surfaces	<i>Panicum capillare</i>	
49 Plants annual, prostrate, mat-forming; blade to 3 cm long, stiff	<i>Munroa squarrosa</i>	
Plants perennial, not prostrate; blade longer than 3 cm	50	
50 Emerging leaves folded	51	
Emerging leaves rolled	53	
51 Sheath margins hyaline; sheath long-pubescent; blades long-pubescent on lower surface, sparsely pubescent above	<i>Danthonia intermedia</i>	
Sheath margins scarios; sheath and blades not as above	52	
52 Plants pilose-pubescent; collar distinct, broad; blade narrow, to 15 cm long	<i>Danthonia unispicata</i>	
Plants sparsely pubescent; collar indistinct, narrow; blade narrow, to 15 cm long, curled	<i>Danthonia spicata</i>	
Plants almost glabrous; collar narrow; blades to 25 cm long	<i>Danthonia parryi</i>	
53 Plants usually decumbent, spreading; blades distinctly veined, tapering to a point ...	<i>Sporobolus cryptandrus</i>	
Plants erect; blades not distinctly veined, bristle-like and often curled	<i>Aristida longiseta</i>	
54 Tips of blades boat-shaped	55	
Tips of blades not boat-shaped	60	
55 Plants annual, low; shoots usually decumbent; blades short, often cross-wrinkled	<i>Poa annua</i>	
Plants perennial, mostly erect	56	
56 Plants short-stoloniferous; blades glossy light green ...	<i>Poa trivialis</i>	
Plants not stoloniferous	57	
57 Plants densely tufted	58	
Plants loosely tufted	59	
58 Shoots very flat, broad; blades flat to folded, erect, to 5 mm wide, light green; ligule to 5 mm long	<i>Helictotrichon hookeri</i>	
Shoots not very flat, small; blades folded, often bristle-like, to 3 mm wide, dark bluish green; ligule to 1 mm long	<i>Poa cusickii</i>	
Shoots not very flat, small; blades flat to folded, to 2 mm wide, green; ligule to 3 mm long	<i>Poa sandbergii</i>	
59 Plants tall, decumbent at base; blades lax, broad at base	<i>Poa palustris</i>	

Plants not tall, not decumbent at base; blades stiff, linear	<i>Poa canbyi</i>
60 Emerging leaves folded	61
Emerging leaves rolled	70
61 Ligule conspicuous, to 10 mm long	62
Ligule inconspicuous, to 1 mm long	63
62 Blades to 10 mm wide, flat, long, indistinctly veined	<i>Dactylis glomerata</i>
Blades to 5 mm wide, flat, long, prominently veined, translucent between veins	<i>Deschampsia caespitosa</i>
Blades to 2 mm wide, short, flexuous, indistinctly veined	<i>Schedonnardus paniculatus</i>
63 Blades flat to folded or involute	64
Blades filiform	68
64 Plants with hard, bulb-like bases; blades tapering to a fine point, dull green; collar glabrous ..	<i>Muhlenbergia cuspidata</i>
Plants without bulb-like bases	65
65 Ligule very short, less than 1 mm long	66
Ligule at least 1 mm long	67
66 Blades short, mostly less than 12 cm long, to 4 mm wide, blue green	<i>Koeleria gracilis</i>
Blades to 50 cm long, to 4 mm wide, gray green	<i>Festuca campestris</i>
67 Ligule about 1 mm long, hyaline to whitish, blunt to rounded	<i>Oryzopsis canadensis</i>
Ligule 1 to 3 mm long, white, usually notched, pubescent	<i>Oryzopsis pungens</i>
68 Plants annual, not densely tufted	<i>Festuca octoflora</i>
Plants perennial, densely tufted	69
69 Plants bluish or gray green; veins indistinct; blades to 25 cm long	<i>Festuca idahoensis</i>
Plants bluish or gray green; veins distinct; blades to 15 cm long	<i>Festuca ovina</i>
Plants gray green; veins distinct; blades to 50 cm long	<i>Festuca campestris</i>
70 Ligule conspicuous, usually more than 3 mm long	71
Ligule inconspicuous, less than 3 mm long	78
71 Plants ascending from spreading base; stoloniferous	<i>Agrostis stolonifera</i>
Plants not stoloniferous	72
72 Sheath margins scarious	73
Sheath margins hyaline	74
73 Plants often densely tufted, erect, bluish green; blades indistinctly veined; saline and alkaline habitats	<i>Puccinellia nuttalliana</i>
Plants not densely tufted, often decumbent, dull green; blades distinctly veined; wet habitats	<i>Alopecurus aequalis</i>
74 Plants annual; blades to 12 mm wide, twisted near tip; dry habitats	<i>Phalaris canariensis</i>

Plants annual; blades to 12 mm wide, not twisted near tip; wet habitats	<i>Beckmannia syzigachne</i>
Plants perennial	75
75 Blades to 12 mm wide, lax; sheath prominently veined, purplish at base	<i>Cinna latifolia</i>
Blades to 5 mm wide	76
76 Plants of moist habitats; blades indistinctly veined, flat, thin	<i>Sphenopholis obtusata</i>
Plants of dry habitats; blades distinctly veined, flat or involute, firm	77
77 Blades usually strongly involute, very long; ligule very conspicuous, to 8 mm long ...	<i>Oryzopsis hymenoides</i>
Blades flat to involute or convolute; ligule conspicuous, obtuse or truncate, to 4 mm long	<i>Stipa comata</i>
Blades flat to involute or convolute; ligule conspicuous, obtuse or truncate, irregular, to 5 mm long	<i>Stipa spartea</i>
78 Plants annual, pubescent; sheath closed to near top	79
Plants perennial, or if annual, not pubescent	80
79 Plants short; blades to 12 cm long, sharp-pointed; collar distinct; sheath pubescent	<i>Bromus tectorum</i>
Plants tall; blades to 30 cm long, tapering, collar indistinct; sheath retrorsely pilose	<i>Bromus marginatus</i>
80 Sheath closed to near top; blades to 10 mm wide, pubescent	<i>Bromus ciliatus</i>
Sheath split, with scarious margins	81
Sheath split, with hyaline margins	83
81 Collar and sheath long-pubescent at margins	<i>Stipa viridula</i>
Collar not pubescent	82
82 Blades to 15 mm wide, bluish green, flat and lax; sheath glabrous	<i>Elymus glaucus</i>
Blades to 10 mm wide, dark green, flat or involute; lower sheath pubescent	<i>Agropyron subsecundum</i>
Blades to 5 mm wide, light green, flat to convolute; sheath glabrous	<i>Agropyron spicatum</i>
83 Blades 10 mm wide or more	84
Blades less than 10 mm wide	85
84 Basal leaves prostrate; collar very narrow, often indistinct, glabrous; blades flat	<i>Oryzopsis asperifolia</i>
Basal leaves erect; collar broad; margins ciliate; base of shoots bulbous; blades twisted	<i>Phleum pratense</i>
Basal leaves erect; collar broad, glabrous; blades not twisted	<i>Elymus virginicus</i>
85 Plants annual; sheath prominently veined, keeled; blades twisted	<i>Lolium persicum</i>
Plants perennial	86
86 Blades soft pubescent, grayish green; collar fairly broad, continuous, pale green	<i>Hordeum jubatum</i>
Blades not pubescent, or only slightly so	87

87 Plants stoloniferous	88
Plants not stoloniferous	89
88 Ligule short, truncate; blades unevenly veined, veins faint at base	<i>Alopecurus pratensis</i>
Ligule long, acute; blades evenly veined, veins equally evident at base	<i>Agrostis stolonifera</i> var. <i>major</i>
89 Blades to 15 cm long; ligule more than 1 mm long	90
Blades more than 15 cm long; ligule to 1 mm long	91
90 Plants glabrous; blades short, usually less than 2 mm wide, sharp-pointed	<i>Agrostis scabra</i>
Plants somewhat pubescent, especially at collar; blades to 5 mm wide, blunt	<i>Trisetum spicatum</i>
91 Sheath somewhat flattened, glabrous; blades linear, bristle-like when dry, glabrous	<i>Stipa richardsonii</i>
Sheath round, lower sheath pubescent; blades flat to involute, scabrous	<i>Agrohordeum macounii</i>
Sheath round, glabrous or softly puberulent; blades flat to involute, finely pubescent	<i>Sitanion hystrix</i>

DESCRIPTIONS OF SPECIES


The order of entry is alphabetical by botanical name. An index to common names begins on page 242.



AGROHORDEUM MACOUNII (Vasey) Lepage

Macoun's wild rye, tufted wild rye



- Growth habit erect, densely tufted, perennial, with fibrous roots
Blade to 5 mm wide, 5 to 20 cm long, erect, rather firm, flat to involute, acuminate, usually scabrous on both sides; midvein extending partway down sheath; rolled at emergence
Sheath round, split, retrorsely pubescent to glabrate, often purplish at base; veins prominent; margins hyaline
Ligule to 1 mm long, membranous, truncate, ciliate
Collar very often indistinct, divided, light or brownish green, glabrous
Auricles absent or rudimentary
Where found throughout area, in moist meadows and borders of woods
Remarks a natural sterile hybrid of *Agropyron trachycaulum* and *Hordeum jubatum*



AGROPYRON CRISTATUM (L.) Gaertn.

crested wheat grass



- Growth habit perennial, tufted, with a dense mass of fibrous roots
- Blade to 8 mm wide, 5 to 20 cm long, flat or slightly involute, acuminate, scabrous and often soft-pubescent on upper surface, smooth or slightly scabrous on lower surface; veins prominent; midvein light-colored; margins scabrous; rolled at emergence
- Sheath round, split, smooth, slightly scabrous or lowest sheaths often soft-pubescent; margins overlapping, hyaline
- Ligule to 1.5 mm long, membranous, truncate, lacerate
- Collar divided, distinct, light or yellowish green, smooth to ciliate
- Auricles to 1.5 mm long, clawlike
- Where found seeded in various soil types because of its drought tolerance
- Remarks introduced; the closely related desert wheat grass, *Agropyron desertorum* (Fisch.) Schutt., has no pubescence on the upper surface of the leaves



AGROPYRON DASYSTACHYUM (Hook.) Scribn.

northern wheat grass



- Growth habit perennial, with rhizomes
Blade to 6 mm wide, 5 to 20 cm long, flat to involute; ridged and very scabrous on upper surface, smooth or sometimes scabrous near base on lower surface; margins scabrous; rolled at emergence
Sheath round, split, smooth or slightly scabrous; midrib often continuing partway down; margins overlapping, scarious
Ligule to 1 mm long, membranous, obtuse, lacerate
Collar divided, smooth, light green
Auricles to 2 mm long, clawlike, clasping, slender
Where found throughout entire area; the most common wheat grass on the Prairies
Remarks awned northern wheat grass, *Agropyron albicans* Scribn. & Smith, a close relative, has narrower leaves



AGROPYRON ELONGATUM (Host) Beauv.

tall wheat grass



Growth habit	tall, erect, loosely tufted, perennial, with fibrous roots
Blade	to 10 mm wide and 20 cm long, appearing glabrous but actually minutely puberulent, acuminate; midrib prominent near collar; rolled at emergence
Sheath	round, split; margins overlapping, hyaline
Ligule	less than 1 mm long, membranous, serrate, often turned downward
Collar	to 2 mm wide, yellow
Auricles	small, clasping, often not seen on older leaves
Where found	hayfields and pastures throughout the area on saline soil
Remarks	introduced by the University of Saskatchewan in 1929 from Siberia; the most salt-tolerant of all cultivated grasses



AGROPYRON INTERMEDIUM (Host) Beauv.

intermediate wheat grass


- Growth habit erect, perennial, with short rhizomes
Blade to 10 mm wide, 5 to 15 cm long, flat, light green, usually glabrous but pubescent in some strains; mid-vein prominent on back; margins ciliate; rolled at emergence
Sheath outer margin usually ciliate, inner margin hyaline
Ligule less than 1 mm long, serrate
Collar to 2 mm wide, yellow, continuous
Auricles variable, slender, clasping, often rudimentary
Where found fairly common in hayfields and pastures in the Parkland
Remarks introduced; often short-lived when grazed; palatable to all classes of livestock; pubescence on blade varies considerably; the introduced pubescent wheat grass, *Agropyron trichophorum* (Link) Richt., has similar vegetative characters



AGROPYRON REPENS (L.) Beauv.

quack grass, couch grass



- Growth habit erect, sod-forming, perennial, with long yellowish white rhizomes
- Blade to 10 mm wide, 6 to 20 cm long, flat or U-shaped, acuminate, slightly keeled at base, green, sometimes glaucous; harsh-scabrous on margins and upper surface, usually sparsely pilose; smooth on lower surface; twisted near tip; rolled at emergence
- Sheath round, split, short, lower sheaths usually pubescent; margins overlapping, hyaline
- Ligule to 1 mm long, membranous, obtuse, entire, lacerate or sometimes ciliate
- Collar divided, distinct, broad, puberulent, yellowish or sometimes tinged with purple
- Auricles to 3 mm long, clawlike, slender
- Where found a weed of waste places and gardens; occasionally in cultivated fields
- Remarks introduced; considerable variation occurs in pubescence of this species; young plants and spring growth are usually more pubescent than older plants or growth produced later in the season. *Agropyron repens* may be distinguished from *Agropyron cristatum* by the puberulent collar and lesser degree of ridging on the leaves. *A. repens* is not tufted, *A. cristatum* is strongly tufted



AGROPYRON SMITHII Rydb.

western wheat grass, bluejoint



- Growth habit erect, sod-forming, perennial, with long slender rhizomes
- Blade to 6 mm wide, 5 to 25 cm long, flat or involute, stiff, acuminate; prominently ridged and scabrous (rarely pubescent) on upper surface, smooth on lower surface; glaucous, bluish green; margins scabrous; blade attached at 45-degree angle to stem; rolled at emergence
- Sheath round, split, prominently veined, glabrous, sometimes brown or purplish at base; margins scarious
- Ligule to 0.5 mm long, membranous, truncate, lacerate, or ciliate
- Collar continuous, medium-broad, not well defined, glabrous, light green, sometimes oblique
- Auricles claw-shaped, clasping, sometimes purplish at base
- Where found throughout area on heavy soils; tolerates drought as well as adequate moisture; alkali-tolerant



AGROPYRON SPICATUM (Pursh) Scribn. & Smith

bluebunch wheat grass



- Growth habit erect, bunch-forming, densely tufted, perennial, with fibrous roots
- Blade to 5 mm wide, 5 to 20 cm long, flat to convolute, ridged; scabrous and hirsute on upper surface, with prominent midrib and veins on lower surface, often minutely pubescent between veins; rolled at emergence
- Sheath round, split, strongly veined, glabrous; margins overlapping, scarious
- Ligule to 0.5 mm long, membranous, truncate, lacerate
- Collar continuous or sometimes divided, glabrous, yellowish
- Auricles to 1.5 mm long; rudimentary to clawlike and fragile if present
- Where found dry prairie in western part of area; common in south central British Columbia



AGROPYRON SUBSECUNDUM (Link) Hitchc.

awned wheat grass



Growth habit	erect, tufted, perennial, with fibrous roots
Blade	to 10 mm wide, 4 to 20 cm long, flat or involute, acuminate; ridged and scabrous on upper surface, slightly scabrous on lower surface; pubescent when young; margins white and very scabrous; rolled at emergence
Sheath	round, split, glabrous in old plants, densely pubescent when young, prominently veined, light green; margins scarious
Ligule	to 1 mm long, membranous, obtuse to truncate, lacerate, brownish
Collar	divided, well defined, light brown
Auricles	absent, or if present, clawlike and often only one
Where found	throughout area, but more common on moist prairie and at forest margins



AGROPYRON TRACHYCAULUM (Link) Malte

slender wheat grass, western rye grass



Growth habit	erect, tufted, perennial, with very short rhizomes
Blade	to 6 mm wide, 5 to 25 cm long, flat, narrowed at base, acuminate, medium green, glaucous; ridged and scabrous on upper surface, keeled and slightly scabrous on lower surface; margins strongly scabrous; rolled at emergence
Sheath	round, split, glabrous, light green, white or purplish at base; margins hyaline
Ligule	to 1 mm long, membranous, truncate, finely ciliate
Collar	continuous, distinct, glabrous, yellowish green, often oblique
Auricles	rudimentary or absent; often only one
Where found	throughout area; the most common wheat grass in situations with good moisture
Remarks	grown for hay for many years before 1940; presently being improved in a plant breeding program; tolerates saline soil



AGROSTIS SCABRA Willd.

hair grass, tickle grass, flyaway grass, rough hair grass


- Growth habit tufted, perennial, with fibrous roots
Blade to 2 mm wide, 2 to 8 cm long, soft, acuminate, often involute or involute and hairlike when dry, glabrous; distinctly ridged and scabrous on upper surface, smooth and distinctly keeled on under surface; margins scabrous; rolled at emergence
Sheath round, slightly keeled, glabrous, smooth, pale green to white or purplish, split; margins overlapping
Ligule to 3 mm long, membranous, truncate or three-pointed, entire or finely lacerate, hairy on outer surface
Collar narrow, inconspicuous, pale green
Auricles absent
Where found throughout area, but common only on moist prairie; one of the first invaders on abandoned farmland



AGROSTIS STOLONIFERA L.

redtop



Growth habit	perennial, with fibrous roots; ascending from a spreading base, the decumbent portion rooting at nodes in wet soil
Blade	to 3 mm wide, flat, deep green; rolled at emergence
Sheath	round, split
Ligule	to 5 mm long, hairy on back, membranous
Collar	continuous, pale green
Auricles	absent or rudimentary
Where found	seeded in lawns, on golf courses, and on landing strips
Remarks	introduced



AGROSTIS STOLONIFERA L. var. *MAJOR* Farw.

creeping bent



Growth habit	stoloniferous, prostrate, perennial, branching and rooting at the nodes, mat-forming
Blade	to 4 mm wide, 3 to 10 cm long, flat, tapering, erect; distinctly ridged on upper surface, slightly keeled on lower surface, scabrous on both surfaces and on margins; rolled at emergence
Sheath	round, split, glabrous, pale green or purplish; margins hyaline
Ligule	to 3 mm long, membranous, thin, entire or finely lacerate, minutely hairy on back
Collar	distinct, smooth, pale green, usually oblique
Auricles	absent
Where found	throughout area in moist places; not common



ALOPECURUS AEQUALIS Sobol.

short-awned foxtail


Growth habit	low, erect, tufted, glabrous, perennial, with fibrous roots
Blade	to 5 mm wide, 4 to 15 cm long, flat, acuminate, dull green, both surfaces and margins scabrous; rolled at emergence
Sheath	round, split, glabrous, green, often purplish at base, with light veins; margins scarios
Ligule	to 5 mm long, membranous, obtuse, minutely pubescent on back, particularly at base
Collar	medium broad, divided, oblique, glabrous, light brown
Auricles	absent
Where found	throughout area around sloughs, springs, and streams; also along roadsides and in shady places in north



ALOPECURUS PRATENSIS L.

meadow foxtail



Growth habit	erect, perennial, forming loose tufts, with very short stolons
Blade	to 6 mm wide, 5 to 15 cm long, flat, acuminate; ridged and scabrous on upper surface, slightly keeled and scaberulous or smooth below; margins very scabrous; rolled at emergence
Sheath	round, split, glabrous, green, often purplish at base, lower sheaths loose, upper ones often inflated; margins overlapping, broad, hyaline
Ligule	to 3 mm long, coarse-membranous, obtuse or truncate, entire, undulate, oblique, faintly striate, ciliate, puberulent on back, variable in shape and margin
Collar	divided, often oblique, glabrous, yellowish
Auricles	absent
Where found	low moist areas and waste places
Remarks	introduced; not common



ANDROPOGON GERARDII Vitman

big bluestem



Growth habit	tall, erect, perennial, with short, thick, scaly rhizomes
Blade	to 10 mm wide, 8 to 50 cm long, flat or V-shaped, constricted at base, acuminate, extending partway down the sheath, slightly scabrous or smooth above, smooth below, with silky hairs at base, light green and often red-tinged, especially near the tip; veins distinct, midvein prominent; margins scabrous; rolled at emergence
Sheath	compressed, split, usually purplish at base and below ground, soft-pubescent or sometimes glabrous; veins distinct; margins hyaline
Ligule	to 3 mm long, membranous, obtuse, ciliate and often lacerate
Collar	medium broad, often indistinct, light or yellowish green, sometimes divided by midvein, pubescent at least on margins
Auricles	absent
Where found	eastern part of area and as far west as the Qu'Appelle Valley; one of the most important species of the true prairie; to 1.2 m tall



ANDROPOGON SCOPARIUS Michx.

little bluestem


- Growth habit erect, tufted, perennial, with short scaly rhizomes
- Blade to 8 mm wide, 5 to 15 cm long, flat to folded, constricted at base; scabrous on upper surface and margins, smooth below; light green, often red-tinged, sometimes glistening, often glaucous; little twisted, tip blunt; folded at emergence
- Sheath compressed, keeled, split, quite open in older leaves, smooth or scabrous, often purplish at base
- Ligule to 2 mm long, membranous, obtuse to truncate, ciliate
- Collar continuous, broad, glabrous, light green or pinkish
- Auricles absent
- Where found throughout area; in Manitoba an important grass of true prairie, westward becoming restricted to areas of adequate moisture or high water table



ARISTIDA LONGISETA Steud.

red three-awn


Growth habit	densely tufted, perennial, with fibrous roots
Blade	to 2 mm wide, 3 to 15 cm long, bristle-like or involute, often curved or flexuous, both surfaces and margins very scabrous upwardly; veins not very distinct; rolled at emergence
Sheath	round, split, prominently veined, having upward-pointing minute prickles, sometimes smooth; margins overlapping, hyaline
Ligule	to 0.5 mm long, a fringe of fine hairs
Collar	indistinct, continuous, retrorsely scabrous, whitish green when evident, with tufts of long hairs at margins
Auricles	absent
Where found	western part of area only; gravelly slopes and dry prairie
Remarks	rare



BECKMANNIA SYZIGACHNE (Steud.) Fern.

slough grass



- Growth habit erect, tufted, annual or biennial, forming large bunches, with fibrous roots
- Blade to 12 mm wide, 8 to 20 cm long, flat, more or less stiff, scabrous above, smooth below; midvein present but not prominent; rolled at emergence
- Sheath round, split, glabrous, smooth; venation muriform; margins overlapping, hyaline
- Ligule to 10 mm long, membranous, obtuse, often lacerate at tip when old
- Collar inconspicuous, divided, smooth, brown or yellowish
- Auricles absent
- Where found in and around shallow sloughs and wet meadows throughout area



BOUTELOUA CURTIPENDULA (Michx.) Torr.

side-oat grama



Growth habit	erect, sod-forming, perennial, with short scaly rhizomes
Blade	to 7 mm wide, 5 to 30 cm long, flat to convolute, acuminate, little twisted, scabrous above, smooth below, with few long hairs scattered on both surfaces; veins distinct; margins scabrous to ciliate; rolled at emergence
Sheath	round, split, usually pubescent, often purplish at the base; veins prominent; margins hyaline
Ligule	to 1 mm long, membranous, truncate, lacerate, ciliate
Collar	medium broad, yellowish green or brown, continuous, with few hairs at the throat
Auricles	absent
Where found	southeastern Saskatchewan along the Souris River and in southwestern Manitoba; rare in Canada



BOUTELOUA GRACILIS (H.B.K.) Lag. ex Steud.

blue grama


- Growth habit densely tufted, perennial, with fibrous roots; occasionally with very short scaly rhizomes
- Blade to 3 mm wide, 2 to 15 cm long, flat to convolute, acuminate, light to medium green, curled; scabrous or pubescent on upper surface especially near base, glabrous on lower surface; veins prominent above and below; rolled at emergence
- Sheath round, split, veins distinct; glabrous or sparsely pilose; margins hyaline
- Ligule to 0.5 mm long; a dense fringe of hairs
- Collar continuous, medium-broad, distinctly yellowish green, with long hairs on inside of margins
- Auricles absent
- Where found throughout area on dry prairie
- Remarks commonly associated with *Stipa comata*; less common northward



BROMUS CILIATUS L.

fringed brome


- Growth habit tall, perennial, with fibrous roots
Blade to 10 mm wide, 8 to 20 cm long, flat, acuminate, dark green, glabrous or soft-pubescent; veins prominent on both surfaces, whitish midvein on lower side extending partway down the sheath; margins scabrous; rolled at emergence
Sheath round, closed to near top, prominently veined, retrorsely pilose or glabrous in older leaves
Ligule to about 1 mm long, coarse-membranous, truncate, entire or lacerate
Collar divided, narrow, glabrous, pale green or brownish; margins sometimes constricted
Auricles absent
Where found throughout the Parkland and the Cypress Hills in fescue prairie and forest margins



BROMUS INERMIS Leys.

smooth brome


- Growth habit tall, perennial, with rhizomes
Blade to 12 mm wide, 15 to 40 cm long, flat, acuminate, dark green, usually glabrous on both surfaces but sometimes minutely pubescent; veins and midrib distinct below; margins scabrous; rolled at emergence
Sheath round, closed to near the top, prominently veined, usually glabrous, sometimes scabrous
Ligule to 1 mm long, membranous, obtuse, entire or lacerate, often brownish at base
Collar narrow, continuous or divided by midvein, glabrous, light or yellowish green
Auricles absent or rarely rudimentary
Where found throughout the entire area, but more commonly in the Parkland
Remarks introduced; the common cultivated brome; known as Austrian brome when introduced into Canada about 1875



BROMUS MARGINATUS Nees

California brome


Growth habit	erect, annual or biennial, with fibrous roots
Blade	to 8 mm wide, to 30 cm long, flat, tapering, prominently veined and pilose on both surfaces; margins scabrous; rolled at emergence
Sheath	slightly compressed to round, split partway, retrorsely pilose, rarely glabrous; veins prominent, midvein continuing down the sheath
Ligule	to 2 mm long, membranous, longitudinally corrugated, obtuse, lacerate
Collar	indistinct, very narrow, divided, brownish green
Auricles	absent
Where found	throughout area in open woods and disturbed places; rare



BROMUS PUMPELLIANUS Scribn.

northern awnless brome



- Growth habit erect, sod-forming, perennial; with strong, creeping, scaly rhizomes
- Blade to 12 mm wide, 10 to 25 cm long, flat, acuminate; veins distinct and very scabrous below, midrib extending partway down sheath; margins scarious and very scabrous; rolled at emergence
- Sheath round, closed to near the top, prominently veined, smooth or scabrous, often purplish at base
- Ligule to 3 mm long, membranous, brown, truncate, lacerate
- Collar continuous, narrow, glabrous, yellowish or light green
- Auricles absent or occasionally rudimentary
- Where found throughout Parkland and Cypress Hills at forest margins and on moist prairie



BROMUS TECTORUM L.

downy brome, downy chess


Growth habit	tufted, annual, with fibrous roots
Blade	to 8 mm wide, 5 to 12 cm long, flat, softly pubescent, sharp-pointed, pale green; margins pilose; rolled at emergence
Sheath	not compressed, keeled, softly pubescent, pale green, often pink- or purple-tinged, split only partway
Ligule	to 3 mm long, membranous, truncate, lacerate
Collar	pubescent, pale, narrow, distinct, usually divided
Auricles	absent
Where found	western part of area; common in British Columbia
Remarks	a troublesome weed in overgrazed pastures and waste places



CALAMAGROSTIS CANADENSIS (Michx.) Beauv.

marsh reed grass



Growth habit	tall, erect, tufted, perennial, with rhizomes
Blade	to 10 mm wide, 7 to 40 cm long, flat, lax, medium green, acuminate; ridged and scabrous above, midrib prominent, smooth or slightly scabrous below; margins scabrous; rolled at emergence
Sheath	round, split, with scarious margins, distinctly veined, glabrous, yellowish at the base
Ligule	to 5 mm long, membranous; truncate, lacerate, often irregular or split
Collar	indistinct, continuous or divided, glabrous, yellowish green, often oblique
Auricles	absent
Where found	throughout entire area in swamps and sloughs



CALAMAGROSTIS INEXPANSA A. Gray

northern reed grass



Growth habit	tall, erect, tufted, perennial, with slender rhizomes
Blade	to 4 mm wide, 10 to 30 cm long, flat or involute, convolute in young shoots, firm; ridged and very scabrous above, smooth or slightly scabrous below; medium to light green; margins scabrous; rolled at emergence
Sheath	round or slightly compressed, split, with scarious overlapping margins, prominently veined, sometimes purplish at base
Ligule	to 6 mm long, membranous, obtuse or truncate, lacerate or slightly ciliate
Collar	indistinct, continuous or divided, glabrous, yellowish green, often oblique
Auricles	absent
Where found	throughout area in wet places and meadows



CALAMAGROSTIS MONTANENSIS Scribn.

plains reed grass



- Growth habit erect, sod-forming, perennial, with slender rhizomes
Blade to 3 mm wide, 5 to 20 cm long, bluish green, stiff, flat to involute, acuminate, veins prominent; ridged and scabrous on the upper surface, faintly midribbed and scabrous below; margins scabrous; rolled at emergence
Sheath round, split, with hyaline margins, prominently veined with midrib extending down the sheath, light green, slightly scabrous
Ligule to 7 mm long, membranous, acute, entire or lacerate in older leaves
Collar indistinct, glabrous, light or brownish green
Auricles absent
Where found throughout northern part of area on dry to moderately moist prairie; occasionally throughout southern Saskatchewan and Alberta



CALAMAGROSTIS NEGLECTA (Ehrh.) Gaertn. Mey. & Schreb.

narrow reed grass


Growth habit	erect, perennial, with long slender rhizomes
Blade	to 5 mm wide, 8 to 30 cm long, flat to convolute, acuminate, ridged and scabrous above, smooth or scabrous below, glaucous, green, usually erect; mid-vein indistinct; margins scabrous; rolled at emergence
Sheath	round, split, with hyaline margins, distinctly veined, glabrous, white or purplish below
Ligule	to 4 mm long, membranous, truncate, lacerate and ciliate
Collar	indistinct, narrow, continuous, glabrous, light or yellowish green
Auricles	absent
Where found	throughout area in wet places
Remarks	usually associated with <i>C. canadensis</i> and <i>C. inexpansa</i> , but much less common



CALAMAGROSTIS RUBESCENS Buckl.

pine grass


- Growth habit erect, sod-forming, perennial, with extensive rhizomes
- Blade to 4 mm wide, 10 to 30 cm long, flat or involute, lax, upper and lower surfaces scaberulous; margins scabrous; rolled at emergence
- Sheath round, split, with scarious margins, distinctly veined, often purplish at base, scabrous
- Ligule to 4 mm long, membranous, obtuse or acute, lacerate when old
- Collar distinct, divided by midrib, pubescent with long soft hairs
- Auricles absent
- Where found western coniferous forests, Cypress Hills, and locally in boreal forests, very common in central British Columbia



CALAMOVILFA LONGIFOLIA (Hook.) Scribn.

sand reed grass



Growth habit	tall, erect, perennial, with long scaly rhizomes
Blade	to 12 mm wide, 10 to 50 cm long, flat to involute, firm, coriaceous, acuminate, smooth on both surfaces, light green; veins not prominent; margins scarious and slightly scabrous; rolled at emergence
Sheath	round, split, distinctly veined, usually smooth but sometimes slightly scabrous or pubescent, light green, often pinkish at base; margins overlapping, hyaline
Ligule	to 3 mm long; a fringe of hairs
Collar	distinct, broad, continuous, yellowish, with tufts of fine hairs on inner margins
Auricles	absent
Where found	throughout entire area on sandy soils
Remarks	one of the most important grasses in sand hills; usually associated with <i>Sporobolus cryptandrus</i> and <i>Oryzopsis hymenoides</i>



CATABROSA AQUATICA (L.) Beauv.

brook grass


- Growth habit perennial, with elongated rhizomes; rooting from prostrate stems
- Blade to 8 mm wide, usually less than 10 cm long, obtuse, flat and flaccid, smooth on both sides, with fine mid-vein; folded at emergence
- Sheath round or slightly compressed, split partway, smooth, and often purplish at base; margins hyaline
- Ligule to 8 mm long, membranous, obtuse
- Collar distinct, narrow, continuous, glabrous, lighter or darker green than leaf blade
- Auricles absent
- Where found throughout area in shallow streams and springs, but not common



CINNA LATIFOLIA (Trev.) Griseb.

slender wood grass


Growth habit	tall, slightly tufted, perennial, with fibrous roots; sometimes bulbous at base
Blade	to 12 mm wide, 10 to 25 cm long, flat, acuminate, finely scabrous on both surfaces; midvein present, extending partway down sheath; rolled at emergence
Sheath	round, split, prominently veined, purplish at base, smooth; margins overlapping, hyaline
Ligule	to 10 mm long, membranous, obtuse, usually lacerate
Collar	divided, distinct, purplish or dark brown, glabrous
Auricles	absent
Where found	throughout area in moist forests



DACTYLIS GLOMERATA L.

orchard grass, cock's-foot



Growth habit	tall, smooth, tufted, perennial, with fibrous roots
Blade	to 10 mm wide, 5 to 25 cm long, flat to folded, tapering, pale green, often glaucous; veins indistinct above and below but with deep furrow over prominent and scabrous midrib; margins scarious, very scabrous; folded at emergence
Sheath	flattened, keeled, split partway, finely but distinctly veined, glabrous, pale green or purplish
Ligule	to 8 mm long, membranous, truncate, and bristle-tipped or obtuse, often split or lacerate, ciliolate
Collar	divided, broad, glabrous, yellowish
Auricles	absent
Where found	seeded in irrigated pastures and orchards; found growing wild in favorable locations
Remarks	introduced



DANTHONIA INTERMEDIA Vasey

wild oat grass, timber oat grass


- Growth habit short, tufted, perennial, with fibrous roots
Blade to 3 mm wide, 5 to 15 cm long, flat to involute, acuminate; somewhat ridged and glabrous or sparsely hairy on upper surface, long-pubescent on lower surface; margins scarios and very slightly scabrous; folded at emergence
Sheath round, split, with hyaline margins, pale green, prominently veined, very pilose-pubescent
Ligule to 1 mm long; a fringe of hairs
Collar distinct, narrow, yellowish green, continuous, with long hairs all around and on inner margins
Auricles absent
Where found Wood Mountain, Cypress Hills, Parkland, foothills of the Rocky Mountains; rare in western Manitoba



DANTHONIA PARRYI Scribn.

Parry oat grass



- Growth habit erect, densely tufted, perennial, with fibrous roots
Blade to 3 mm wide, 10 to 25 cm long, flat to involute, mostly erect-flexuous and narrow or filiform, scaberulous on upper surface and on the scarious margins, glabrous below, yellowish green; veins distinct above but no midrib evident; folded at emergence
Sheath round, split, distinctly ridged at the upper end, yellowish and glabrous; margins overlapping, scarious
Ligule to 0.5 mm long; a fringe of hairs
Collar a narrow dark continuous ridge, with tufts of long stiff hairs at margins
Auricles absent
Where found foothills of the Rocky Mountains as far north as Rocky Mountain House, rarer north of the Bow River
Remarks blades break off at the collar during the winter



DANTHONIA SPICATA (L.) Beauv. ex Roem. & Schult.

poverty oat grass



Growth habit	tufted, perennial, with fibrous roots
Blade	to 3 mm wide, basal leaves 5 to 15 cm long, flat to convolute, acuminate, flexuous, usually sparsely pilose but often glabrous; prominently veined on both surfaces, under surface bright green and sometimes glossy, upper surface dull green or glaucous; margins scabrous; folded and involute at emergence
Sheath	round, split, with scarious margins, pale green to white, usually pilose-pubescent
Ligule	to 1.2 mm long; a fringe of hairs
Collar	narrow, continuous, often indistinct, light or yellowish green with long hairs at margins
Auricles	absent
Where found	throughout area, in moderately dry prairie but not common; occasionally abundant



DANTHONIA UNISPICATA (Thurb.) Munro ex Macoun

one-spike oat grass, dwarf oat grass



- Growth habit short, erect, tufted, perennial, with fibrous roots
Blade to 3 mm wide, 5 to 15 cm long, flat to involute, light green, pilose above, glabrous or slightly pilose below, light green, acuminate; margins scariosus and scabrous; veins not very prominent; folded at emergence
Sheath round, split, prominently veined; glabrous or upper sheaths pilose; margins overlapping, scariosus
Ligule a fringe of fine hairs
Collar distinct, broad, flared up the blade, continuous, yellowish, with tufts of long hairs at margins
Auricles absent
Where found western part of area, in moderately dry prairie; not common



DESCHAMPSIA CAESPITOSA (L.) Beauv.

tufted hair grass


- Growth habit erect, densely tufted, perennial, with fibrous roots
- Blade to 5 mm wide, 5 to 30 cm long, flat to folded, rather firm and bristle-like, contracted at collar, often flexuous, prominently ridged and scabrous above, smooth or slightly scabrous below; translucent between veins, light green; margins scabrous; folded at emergence
- Sheath compressed, keeled, prominently veined, split, light green or straw-colored; margins overlapping, hyaline
- Ligule to 10 mm long, membranous, acuminate, entire or sometimes split, very prominent, continuous with margins of sheath
- Collar narrow, divided, sometimes indistinct, yellowish green or brownish, a noticeable swelling occurring at each end of collar
- Auricles absent
- Where found throughout area in moist places; an important grass of moist meadows in the foothills of the Rocky Mountains, and northward beyond Peace River



DISTICHLIS STRICTA (Torr.) Rydb.

desert salt grass, alkali grass, salt grass


- Growth habit sod-forming, perennial, with extensive scaly rhizomes
Blade to 4 mm wide, 1 to 15 cm long, acuminate, flat to involute, ridged, scabrous on upper surface with a few long hairs especially near base, smooth below; margins very scabrous; lower blades much shorter than upper ones; folded at emergence
Sheath round, split, usually glabrous, basal ones usually yellowish; margins overlapping, scarious
Ligule a fringe of hairs to 0.5 mm long; fused at the base
Collar narrow, continuous, distinct, yellowish, with tufts of long hairs at ends
Auricles absent
Where found throughout area; common in saline or alkaline areas, occasionally on dry slopes



ECHINOCHLOA CRUSGALLI (L.) Beauv.

barnyard grass



Growth habit	annual, tufted, with fibrous roots
Blade	to 15 mm wide, 10 to 30 cm long, flat or V-shaped, glabrous, pale or yellowish green, keeled below; margins smooth or scabrous; rolled at emergence
Sheath	compressed, keeled, split, smooth, glabrous, pale green; margins hyaline
Ligule	absent
Collar	broad, continuous, glabrous, yellowish green
Auricles	absent
Where found	in gardens and waste places
Remarks	a weed in wheat fields and gardens



ELYMUS ANGUSTUS Trin.

Altai wild rye


- Growth habit tufted, perennial, with fibrous roots
Blade to 15 mm wide, 5 to 40 cm long, stiff, flat, acuminate, prominently veined; retrorsely scabrous on upper surface, smooth below; stomata appearing as white lines on under surface; margins retrorsely scabrous; rolled at emergence
Sheath split, prominently veined, glabrous; margins hyaline
Ligule to 2 mm long, membranous, acute, lacerate, ciliate
Collar broad, continuous, yellowish, glabrous, constricted
Auricles to 3 mm long, prominent, clawlike, clasping, hyaline
Where found in hay and pasture fields
Remarks introduced from Mongolia; a very coarse grass that is eaten readily by cattle in the late summer and autumn



ELYMUS CANADENSIS L.

Canada wild rye, nodding wild rye



- Growth habit tall, erect, perennial, with short rhizomes
- Blade to 20 mm wide, 5 to 40 cm long, flat to convolute, acuminate, prominently veined above and below, dark green, sometimes glaucous; upper surface dull and slightly scabrous, lower surface glabrous with prominent midrib; margins scabrous; rolled at emergence
- Sheath round, split, prominently veined, green or glaucous, base often purplish; margins overlapping, scarios, the inner usually broadly hyaline, the outer often ciliate
- Ligule to 2 mm long, coarse-membranous, obtuse or truncate, lacerate and short-ciliate
- Collar broad, continuous, often oblique, glabrous, yellowish or light green
- Auricles to 2 mm long, clawlike, clasping
- Where found throughout area; in sand hills, along stream banks and in woods
- Remarks makes poor bedding; seldom eaten by livestock



ELYMUS CINEREUS Scribn. & Merr.

giant wild rye


- Growth habit very tall, tufted, perennial, with short thick knotted rhizomes
- Blade to 15 mm wide, 15 to 50 cm long, flat to convolute, acuminate, prominently veined and scabrous above, scaberulous and less distinctly veined below, fairly thick and stiff; margins scarios and often scabrous; rolled at emergence
- Sheath round, split, prominently but finely veined, smooth or scaberulous, light green, base brownish or faintly purple; margins scarios
- Ligule to 7 mm long, membranous, obtuse or truncate, lacerate
- Collar narrow, continuous, yellow, sometimes divided and indistinct; glabrous
- Auricles to 4 mm long, clawlike, clasping
- Where found along riverbanks, ravines, and moist slopes; uncommon; occasionally seeded
- Remarks single plants to 2 m tall, and often over 50 cm wide at the base



ELYMUS GLAUCUS Buckl.

blue wild rye


Growth habit	tall, tufted, perennial, with fibrous roots
Blade	to 15 mm wide, 5 to 30 cm long, flat, becoming convolute on drying, usually lax, tapering but lower blades not sharp-pointed, scabrous on both surfaces, not prominently veined; margins scarious and scabrous; rolled at emergence
Sheath	round, split, smooth or scabrous, distinctly veined; margins overlapping, scarious
Ligule	to 2 mm long, membranous, truncate, lacerate
Collar	narrow, continuous or sometimes divided, glabrous, yellowish or brownish, coriaceous at the margins, often oblique
Auricles	to 4 mm long, clawlike, clasping, sometimes absent
Where found	throughout area in moderately moist, open woods and meadows; not common



ELYMUS INNOVATUS Beal

hairy wild rye



Growth habit	tall, erect, perennial, with slender scaly rhizomes
Blade	to 12 mm wide, 5 to 25 cm long, flat to convolute, somewhat rigid, acuminate, scabrous above and below; margins very scabrous; veins more prominent above than below; rolled at emergence
Sheath	round, split, lower sheaths scabrous, upper sheaths usually smooth, prominently veined; margins scarios
Ligule	to 0.5 mm long, membranous, truncate
Collar	indistinct, narrow, divided, light green
Auricles	to 2 mm long, clawlike
Where found	throughout area in open woods and clearings; rare in the south but often abundant in the north
Remarks	seldom eaten by cattle



ELYMUS JUNCEUS Fisch.

Russian wild rye



- Growth habit deep-rooted, tufted, perennial, with fibrous roots
Blade to 5 mm wide, 5 to 25 cm long, flat to convolute, acuminate, erect, somewhat scabrous on both surfaces, prominently veined; margins scabrous; rolled at emergence
Sheath round, split, prominently veined and smooth; margins overlapping, hyaline
Ligule to 1 mm long, membranous, obtuse to truncate, lacinate, ciliate
Collar broad, continuous, glabrous, yellowish or light green
Auricles to 2 mm long, clawlike, clasping
Where found seeded for pasture in the prairie area of Western Canada
Remarks introduced from Siberia; becoming the most common seeded grass in the southern prairies; slow to establish but very persistent



ELYMUS VIRGINICUS L.

Virginia wild rye, lyme grass



- Growth habit tall, loosely tufted, perennial, with fibrous roots
Blade to 12 mm wide, 10 to 30 cm long, flat, acuminate, scabrous on both surfaces, distinctly nerved, somewhat keeled below; margins scabrous; rolled at emergence
Sheath round, split, smooth, scabrous or sparsely retrorsely pubescent, light green, sometimes purplish at the base; margins overlapping, hyaline, the outer ciliate
Ligule 0.5 mm long, thick-membranous, greenish or brown-tinged, truncate, wavy, ciliate
Collar broad, continuous, glabrous, yellowish, often oblique
Auricles to 1.5 mm long, clawlike, sharp, occasionally absent
Where found throughout area in woods, clearings, and moist meadows
Remarks this species may be distinguished from *Elymus canadensis* by its narrower blades, shorter ligule, and more scabrous blade surfaces



FESTUCA CAMPESTRIS Rydb.

rough fescue


- Growth habit erect, densely tufted, perennial, with fibrous roots, occasionally with short rhizomes
- Blade to 4 mm wide, to 50 cm long, acuminate, flat or folded; ridged and scabrous on the inner surface, scabrous on outer surface; margins scabrous; folded at emergence
- Sheath round or slightly compressed, smooth or scabrous; split, margins hyaline, overlapping
- Ligule to 0.5 mm long, membranous, truncate, lacerate, ciliate
- Collar very narrow, distinct, glabrous, yellowish
- Auricles rudimentary
- Where found foothills of the Rocky Mountains and western Cypress Hills
- Remarks single plants are often 50 cm or more across when protected from grazing; plants in the Cypress Hills often have short rhizomes and may be crosses with plains rough fescue



FESTUCA ELATIOR L. var. *PRATENSIS* A. Gray

meadow fescue


- Growth habit tufted, perennial, with rhizomes
- Blade to 8 mm wide, 10 to 50 cm long, bright green; upper surface dull, scabrous, and prominently veined, lower surface glossy, smooth, slightly keeled; margins scabrous; rolled at emergence
- Sheath oval, glabrous, pale green, reddish to purple at base, split to near base; margins overlapping, hyaline
- Ligule to 0.5 mm long, membranous, greenish, truncate to obtuse, entire
- Collar broad, distinct, glabrous, yellow green to cream-colored, divided by midrib; margins thin, dilated, often flexuous
- Auricles to 1.5 mm long, present, soft, clawlike or blunt, yellow green to creamy white
- Where found occasionally in cultivated hay meadows and pastures in Manitoba and western Alberta; not common
- Remarks introduced from Europe in seed mixtures; reed or tall fescue, *Festuca elatior* L. var. *arundinacea* (Schreb.) Wimm., is coarser and has scabrous leaves; it is sown in wetter places than meadow fescue



FESTUCA HALLII (Vasey) Piper

plains rough fescue


- Growth habit erect, densely tufted, perennial, often with short rhizomes
Blade to 1.5 mm wide, to 30 cm long, acuminate, permanently folded; ridged and scabrous on the inner surface, scabrous on the outer surface; margins scabrous; folded at emergence
Sheath round, ridged, split, scabrous, margins scarious
Ligule a lacerate, ciliate fringe
Collar indistinct, yellowish
Auricles absent
Where found Parklands throughout the area; common in Wood Mountain and eastern Cypress Hills; occasionally on north-facing slopes in the plains
Remarks single plants are seldom more than 10 cm across when protected from grazing



FESTUCA IDAHOENSIS Elmer

bluebunch fescue


Growth habit	erect, densely tufted, perennial, with fibrous roots
Blade	to 2 mm wide, 5 to 25 cm long, folded, filiform, scabrous, bluish green; veins indistinct; blades so erect as to appear continuous with sheath; folded at emergence
Sheath	flattened, keeled, split only partway, smooth or scabrous, basal sheaths pinkish
Ligule	to 1 mm, truncate, ciliate
Collar	indistinct
Auricles	absent
Where found	Cypress Hills and southern foothills of Rocky Mountains; not common but locally abundant



FESTUCA OCTOFLORA Walt.

six-weeks fescue


Growth habit	low, erect, tufted, annual, with fibrous roots
Blade	to 2 mm wide, 2 to 10 cm long, filiform, twisted, prominently ridged and usually scaberulous below, dark green; folded at emergence
Sheath	flattened, keeled, ridged, usually retrorsely scabrous or minutely pubescent, split; margins overlapping, scarious
Ligule	to 1 mm long, membranous, truncate, ciliate, lacerate
Collar	continuous, narrow, distinct, glabrous, light green
Auricles	absent
Where found	throughout area on moist open ground, and in waste places; not common



FESTUCA OVINA L.

sheep fescue



Growth habit densely tufted, perennial, with fibrous roots
Blade to 1 mm wide, 5 to 15 cm long, filiform, permanently folded, bluish green, glaucous, ridged; scabrous on inner surface, usually scabrous on outer surface; margins smooth; folded at emergence
Sheath round or slightly compressed, split, glabrous or finely pubescent, bluish green, glaucous; margins overlapping, broad, hyaline
Ligule membranous, truncate, very short or rudimentary, ciliate
Collar indistinct, narrow, glabrous, yellowish
Auricles absent
Where found throughout area in moist prairie on various soil types
Remarks *F. ovina* var. *ovina* is naturalized from Eurasia and occurs in the plains; the native *F. ovina* var. *saximontana* (Rydb.) Gleason occurs in the Rocky Mountains; it is distinguished by floral characters and leaf anatomy.



FESTUCA RUBRA L.

red fescue


- Growth habit tufted, erect, perennial, with short rhizomes
- Blade to 3 mm wide, 5 to 15 cm long, thick, V-shaped to closely folded, dark green; deeply ridged on upper surface, smooth and slightly shiny on lower surface; margins smooth; folded at emergence
- Sheath round or oval, finely pubescent, split partway only; dead basal sheaths reddish brown, often chartaceous
- Ligule to 0.5 mm long, membranous, truncate, entire or ciliate
- Collar indistinct, narrow, continuous, glabrous, pale green
- Auricles absent, or present as rounded extensions of ends of collar
- Where found in the foothills of the Rocky Mountains and northern Alberta as a native species
- Remarks commonly cultivated for pasture, hay, and lawns; the cultivated variety was selected from importations from the Atlas Mountains in Africa



GLYCERIA GRANDIS S. Wats.

tall manna grass


- Growth habit tall, stout, glabrous, perennial, with rhizomes
- Blade to 15 mm wide, 10 to 40 cm long, flat or V-shaped, tapering to a boat-shaped tip, glabrous above and below, slightly scabrous on margins, prominently but finely veined below, with white midvein; muriform; folded at emergence
- Sheath compressed and keeled, closed to near top, but rupturing easily; smooth or slightly scabrous, pale or yellowish green, strongly striate and crossveined
- Ligule to 4 mm long, membranous, truncate to abruptly acuminate, entire or slightly undulate
- Collar conspicuous, divided, pale green or yellowish brown
- Auricles absent
- Where found throughout area in sloughs, along streams, and in wet meadows
- Remarks plants are yellowish green, coarse, with rather firm leaves having crossveins on both blade and sheath; seeds are small and very hard



GLYCERIA STRIATA (Lam.) Hitchc.

fowl manna grass


- Growth habit erect, loosely tufted, perennial, with short rhizomes
- Blade to 6 mm wide, 6 to 30 cm long, flat or V-shaped, tapering to a sharp boat-shaped tip, pale green, glabrous above and below, scabrous on margins, thin, soft, crossveined; folded at emergence
- Sheath slightly flattened, somewhat keeled, closed to near top, smooth or slightly scabrous, pale green or purplish-tinged at base, finely nerved and crossveined
- Ligule to 5 mm long, membranous, obtuse, entire or lacerate
- Collar divided or continuous, indistinct, pale green, glabrous
- Auricles absent
- Where found throughout area in sloughs, along streams, and in wet meadows
- Remarks this species is distinguished from *G. grandis* by its narrower leaves, purplish-tinged sheaths, and less conspicuous crossveins on blades



HELICTOTRICHON HOOKERI (Scribn.) Henr.

Hooker's oat grass


- Growth habit erect, densely tufted, smooth-leaved, perennial, with fibrous roots
- Blade to 5 mm wide, 2 to 12 cm long, flat to folded, firm, very erect, linear and boat-shaped at the tip, smooth; midrib prominent and thickened; margins scabrous, whitish; folded at emergence
- Sheath compressed, keeled, split, prominently veined, glabrous, rather short, pale green at base; margins hyaline
- Ligule to 5 mm long, membranous, obtuse or acute
- Collar divided, often inconspicuous, yellowish, light brown, or purplish
- Auricles absent
- Where found throughout area in places of moderate moisture; seldom abundant



HIEROCHLOE ODORATA (L.) Beauv.

sweet grass


- Growth habit erect, sweet-smelling, sod-forming, perennial, with extensive rhizomes
- Blade to 8 mm wide, 2 to 20 cm long, flat to convolute, tapering to an obtuse point, scabrous or smooth on either surface, dark green, with fine prominent nerves on both sides; midrib white below; margins scabrous; rolled at emergence
- Sheath round, split, prominently veined, smooth or slightly scabrous, young shoots purplish at base; margins scarious
- Ligule to 3 mm long, membranous, obtuse, toothed, lacerate, sometimes ciliate
- Collar narrow, divided, light green, glabrous
- Auricles absent
- Where found throughout area in moist prairie and marsh; often an early invader on abandoned land



HORDEUM JUBATUM L.

foxtail barley, wild barley


Growth habit	erect, densely tufted, perennial, with fibrous roots
Blade	to 6 mm wide, 5 to 15 cm long, flat or U-shaped, acuminate, twisted, prominently veined, scabrous and usually villose above, densely puberulent or scabrous below, rarely smooth, bluish green, often purplish; margins scabrous; rolled at emergence
Sheath	round, split, distinctly veined, glaucous, pubescent, often purplish; margins overlapping, hyaline
Ligule	to 1 mm long, membranous, obtuse or truncate, lacerate, finely ciliate
Collar	medium broad, continuous, glabrous, pale green, often purplish, sometimes oblique
Auricles	absent or rudimentary
Where found	throughout area in waste places; around sloughs and marshes, and along streams; tolerates moderate alkalinity and flooding



KOELERIA GRACILIS Pers.

June grass



- Growth habit low, erect, tufted, perennial, with fibrous roots
- Blade to 4 mm wide, 5 to 12 cm long, tapering to an obtuse tip, flat to involute, bluish green, stiff, ridged and scabrous above, keeled and usually smooth underneath, sometimes glabrous or pubescent on both surfaces; margins narrowly scarious and scabrous; folded at emergence
- Sheath round, split, scabrous and often pubescent, distinctly veined, pale green; margins scarious
- Ligule to 1 mm long, membranous, truncate to obtuse, often split, usually ciliate
- Collar medium broad, continuous or divided, yellowish green, smooth or hairy on margins
- Auricles absent
- Where found throughout area; one of the most common and widespread species, but rarely abundant
- Remarks has different growth forms in different environments, with short basal leaves in dry prairie but with few basal leaves and a taller growth where more moisture is available



LOLIUM PERENNE L.

perennial rye grass



- Growth habit tufted, perennial, with fibrous roots
Blade to 6 mm wide, 5 to 15 cm long, acuminate, keeled, prominently ridged on upper surface, smooth and glossy on lower surface, bright green; margins slightly scabrous; folded at emergence
Sheath usually compressed but sometimes almost cylindrical, not keeled, glabrous, pale green, reddish at base, closed or split
Ligule to 2 mm long, thin-membranous, obtuse, toothed near apex
Collar narrow, distinct, glabrous, yellowish to whitish green
Auricles small, soft, clawlike
Where found seeded in short-term pasture and hayland, and in lawn-grass mixtures
Remarks this introduced species resembles *Festuca elatior* except that it has leaves folded at emergence; a common pasture grass in western Europe, New Zealand, and northeastern United States



LOLIUM PERSICUM Boiss. & Hohen.

Persian darnel



Growth habit	erect, annual, with fibrous roots
Blade	to 6 mm wide, 5 to 15 cm long, flat to convolute, acuminate, twisted, ridged and scabrous above, distinctly veined and smooth or slightly scabrous below; margins slightly scabrous; rolled at emergence
Sheath	round or slightly flattened, split, prominently veined, keeled, glabrous; margins overlapping, hyaline
Ligule	to 1 mm long, membranous, brownish, truncate to rounded, lacerate
Collar	often indistinct, divided, sometimes oblique, yellowish or brown
Auricles	usually absent, occasionally rudimentary
Where found	in grain fields, gardens, and waste places
Remarks	introduced; a troublesome weed in grain fields; it will compete with wheat to a point where no grain will be harvested



MUHLENBERGIA ASPERIFOLIA (Nees & Mey.) Parodi

scratch grass



- Growth habit low, erect, perennial, with slender scaly elongated rhizomes
- Blade to 2 mm wide, 2 to 5 cm long, flat, upper surface very scabrous; lower surface smooth, midrib prominent; margins scabrous; folded at emergence
- Sheath flattened, keeled, split, glabrous, medium green, prominently veined; margins hyaline
- Ligule to 1 mm long, membranous, lacerate, truncate
- Collar distinct, glabrous, divided by prominent midvein part-way down the sheath
- Auricles absent
- Where found throughout area in moist prairie and meadows; moderately alkali-tolerant; not common



MUHLENBERGIA CUSPIDATA (Torr.) Rydb.

prairie muhly



- Growth habit erect, densely tufted, perennial, with hard, bulblike scaly bases, with fibrous roots
- Blade to 3 mm wide, 3 to 10 cm long, flat to folded, acuminate, prominently veined and finely rough hairy above, glabrous with prominent midvein below; margins scabrous and scarious; folded at emergence
- Sheath slightly flattened, very short, split, glabrous or slightly puberulent; margins overlapping
- Ligule to 1 mm long, membranous, truncate or rounded, lacinate, ciliate
- Collar broad, continuous, light yellow, glabrous
- Auricles absent
- Where found throughout area on very dry prairie, becoming rare towards the Parklands; often abundant on eroded calcareous slopes



MUHLENBERGIA RACEMOSA (Michx.) B.S.P.

marsh muhly



Growth habit	erect, perennial, with stout scaly rhizomes
Blade	to 6 mm wide, 5 to 15 cm long, flat to folded, abruptly acuminate, dark green, scabrous on both surfaces; veins not prominent, midvein prominent below; margins scarious and scabrous; folded at emergence
Sheath	flattened, keeled, split, distinctly veined, smooth or scabrous; margins hyaline
Ligule	to 1 mm long, membranous, truncate to rounded, lacerate
Collar	continuous, narrow, yellowish or brownish green, glabrous
Auricles	absent
Where found	throughout area in moist meadows and margins of woods
Remarks	<i>M. andina</i> (Nutt.) Hitchc. has been found in western Manitoba. It differs in having a ligule acute to 3 mm long.



MUHLENBERGIA RICHARDSONIS (Trin.) Rydb.

mat muhly



- Growth habit low, erect, densely tufted, perennial, with extensive scaly rhizomes
- Blade to 2 mm wide, to 10 cm long, flat or involute; upper surface finely striate, slightly scabrous; lower surface smooth, midrib prominent; margins scabrous; folded at emergence
- Sheath round, split, smooth, medium green, margins hyaline
- Ligule to 3 mm long, membranous, obtuse to acute, entire, sometimes lacerate at tip
- Collar to 1 mm wide, distinct, glabrous, greenish white
- Auricles absent
- Where found throughout area on moist prairie; tolerates slight alkalinity



MUNROA SQUARROSA (Nutt.) Torr.

false buffalo grass

Growth habit	low, tufted, annual, prostrate, mat-forming, with fibrous roots
Blade	to 3 mm wide, 1 to 3 cm long, crowded at the nodes and ends of the branches, stiff, spreading, scabrous, flat sharp-pointed, in fascicles; rolled at emergence
Sheath	pilose at throat, often ciliate, margins loose, soon becoming papery, inflated
Ligule	to 1 mm long, a fringe of hairs
Collar	divided, narrow
Auricles	absent
Where found	extreme southern parts of area on dry prairie; rare, known from only a few locations
Remarks	introduced from the United States





H. F. Best

ORYZOPSIS ASPERIFOLIA Michx.

white-grained mountain rice grass


Growth habit	erect, tufted, perennial, with short rhizomes
Blade	of two types, some only 1 cm long, others to 40 cm long, all 3 to 10 mm wide, flat to convolute, erect, tapering at each end; surfaces smooth or slightly scabrous, distinctly veined above; margins very scabrous; rolled at emergence
Sheath	round, rather short, split, smooth or scaberulous, dark purple at base; margins slightly overlapping, hyaline
Ligule	to 1 mm long, membranous, ciliate, lacerate
Collar	very narrow, continuous, light green, often indistinct
Auricles	absent
Where found	throughout the Cypress Hills, the Parkland, and adjacent forest



ORYZOPSIS CANADENSIS (Poir.) Torr.

Canadian rice grass



- Growth habit erect, densely tufted, perennial, with fibrous roots
Blade to 4 mm wide, to 30 cm long, flat to involute, strongly ridged, scabrous
Sheath round, split, the margins scarious
Ligule to 1.5 mm long, hyaline to whitish or greenish, blunt to rounded
Collar narrow, indistinct, yellowish
Auricles absent
Where found northern parklands, Boreal forest, Riding Mountain, grassy openings; not common
Remarks can be mistaken for plains rough fescue or Richardson's needle grass; the ligule and usually flatter or more loosely folded leaf are distinguishing marks



ORYZOPSIS HYMENOIDES (Rœm. & Schult.) Ricker

Indian rice grass


Growth habit	tall, tufted, perennial, with fibrous roots
Blade	to 5 mm wide, to 50 cm long, involute, seldom flat, long-tapering, coriaceous, coarsely ridged and scabrous above, prominently veined and smooth below; rolled at emergence
Sheath	round, split, very prominently veined, smooth or slightly scabrous; margins overlapping, hyaline, outer often fringed with hairs
Ligule	to 8 mm long, membranous, acute, often split, very conspicuous
Collar	indistinct, narrow, yellowish, often with small tufts of hairs at the margins
Auricles	absent
Where found	throughout area on sandy soils
Remarks	an important grass in sand hills, very resistant to wind action and grazing; palatable to livestock



ORYZOPSIS PUNGENS (Torr.) Hitchc.

northern rice grass



Growth habit	erect, densely to loosely tufted, perennial, with fibrous roots
Blade	to 3 mm wide, to 20 cm long, flat to involute, strongly ridged, dark green, scabrous
Sheath	round, strongly ribbed, scabrous, split; margins overlapping, hyaline to purplish
Ligule	to 2 mm long, thick, whitish or yellowish, noded, ribbed, rounded, pubescent
Collar	indistinct, narrow, yellowish
Auricles	absent
Where found	open woods, clearings, and grassy areas in Boreal forest
Remarks	somewhat similar in appearance to plains rough fescue, but dark green leaves and ligule are distinguishing marks



PANICUM CAPILLARE L.

witch grass

- Growth habit erect, tufted, annual, with fibrous roots
Blade to 15 mm wide, 8 to 25 cm long, hispid on both surfaces, distinctly veined below, less so above, flat, acuminate, soft, keeled at base; margins scabrous, papillose-ciliate toward base; rolled at emergence
Sheath round, split, prominently veined, dull green, often purple-tinged, conspicuously papillose-hispid; margins overlapping, hyaline
Ligule to 2 mm long, a dense fringe of hairs fused at base
Collar usually broad, continuous, often indistinct, yellowish green, pubescent
Auricles absent
Where found eastern part of area on light soils and in waste places




K.A. BENT

PANICUM VIRGATUM L.

switch grass


- Growth habit erect, perennial, with large scaly rhizomes
Blade to 12 mm wide, 10 to 50 cm long, flat, acuminate, constricted at base, distinctly veined, with prominent midvein, usually glabrous on both surfaces but long-pubescent on upper surface at base; margins scarios and scabrous; rolled at emergence
Sheath round, split, with scarios and villose margins, glabrous, prominently veined, white to purplish-tinged below
Ligule to 3 mm long, a fringe of hairs fused at the base
Collar broad, continuous, yellowish, pubescent on margins
Auricles absent
Where found eastern part of area on moist prairie; rare



PHALARIS ARUNDINACEA L.

reed canary grass



- Growth habit tall, stout, erect, perennial, with long scaly pinkish rhizomes
- Blade to 15 mm wide, 10 to 30 cm long, flat, acuminate, light green, glaucous, finely veined, smooth or slightly scabrous on both surfaces, midrib prominent below; margins scabrous; rolled at emergence
- Sheath round, split, glabrous, light or yellowish green, distinctly veined, pinkish at base; margins overlapping, hyaline
- Ligule to 5 mm long, membranous, white, minutely pubescent on back, acute or obtuse, entire, lacerate or split
- Collar divided, or narrowed at midrib if continuous, glabrous, yellowish green, usually oblique
- Auricles absent
- Where found throughout area in wet places
- Remarks imported from Sweden for forage in periodically flooded areas and on irrigated soils; native forms are known



PHALARIS CANARIENSIS L.

Canary grass



Growth habit	erect, annual, with fibrous roots
Blade	to 12 mm wide, 5 to 30 cm long, flat, acuminate, twisted near apex, scabrous on both surfaces, finely nerved, midrib prominent below and extending part-way down sheath; margins scarios and slightly scabrous; rolled at emergence
Sheath	round, or flattened, split, prominently veined, scabrous; margins overlapping, hyaline
Ligule	to 8 mm long, membranous, acute, lacerate
Collar	narrow, continuous, often indistinct, light or yellowish green, wider at ends than at center
Auricles	absent
Where found	in gardens and waste places, dry habitats
Remarks	introduced; grown for birdseed



PHLEUM ALPINUM L.

alpine timothy


- Growth habit erect or decumbent at base, densely tufted, perennial, with a few rhizomes
- Blade to 8 mm wide, 2 to 15 cm long, flat, tapering, usually short, finely veined above, with prominent midvein below; scabrous on upper surface and margins, smooth or scaberulous below; rolled at emergence
- Sheath round, split, inflated near middle, prominently veined, glabrous, light green, often purplish at base; margins overlapping, hyaline
- Ligule to 4 mm long, membranous, truncate or obtuse, often notched on one side, rarely lacerate
- Collar narrow, often oblique, glabrous, continuous or divided, distinct, light or yellowish green
- Auricles absent
- Where found Cypress Hills, foothills of the Rocky Mountains, and Parkland; moist meadows



PHLEUM PRATENSE L.

timothy


- Growth habit erect, tufted, perennial, each stem growing from a bulbous or swollen base, with fibrous roots
- Blade to 12 mm wide, 5 to 25 cm long, flat, acuminate, distinctly veined, smooth or scaberulous on both surfaces, twisted, erect, light green, glaucous; scarious margins scabrous, retrorsely so at base; rolled at emergence
- Sheath round, split, distinctly veined, glabrous, light green, often purplish at base; margins overlapping, hyaline
- Ligule to 3 mm long, membranous, usually obtuse
- Collar medium broad, continuous, light or yellowish green, sometimes oblique, margins sparsely retrorse-ciliate
- Auricles absent
- Where found commonly seeded for hay and pasture in foothills and Parklands
- Remarks introduced into the United States from Europe about 1700; named for Timothy Hanson, an eighteenth-century American farmer



PHRAGMITES COMMUNIS Trin.

common reed, cane grass


- Growth habit very stout, erect, perennial, with stout creeping rhizomes
- Blade to 30 mm wide, 15 to 40 cm long, acuminate, narrowed at base, glabrous above and below; margins scarios and scabrous; veins fine, not very prominent; rolled at emergence
- Sheath round, split, margins scarios, glabrous, not prominently veined, light yellowish green, purplish at base
- Ligule to 1 mm long, a fringe of hairs, thick, brown, truncate, lacerate, long ciliate, pubescent on back
- Collar broad, continuous, glabrous, yellowish and usually brown on the margins
- Auricles absent or rudimentary
- Where found throughout area in wet places; fairly common in Qu'Appelle Valley
- Remarks the tallest of our native grasses, growing to 3 m tall



POA ANNUA L.

annual blue grass, low spear grass, annual meadow grass



- Growth habit low-growing, tufted, annual, with fibrous roots
Blade to 4 mm wide, 2 to 8 cm long, flat or V-shaped at base, with boat-shaped tip, light green, not glossy, thin, soft, often crosswrinkled; having two distinct light lines along midvein; margins glabrous, slightly scabrous toward tip; blades widely spreading from axis of shoot; folded at emergence
Sheath compressed and slightly keeled, glabrous, light green, split partway only; margins usually overlapping, membranous
Ligule to 3.0 mm (usually 1.2 to 1.8 mm) long, membranous, white, acute, entire
Collar distinct, glabrous, pale green, V-shaped
Auricles absent
Where found throughout area in waste places, gardens, lawns, barnyards, and footpaths
Remarks its tufted growth, its soft, pale green and generally puckered blades, and its white conspicuous ligule are diagnostic; often found growing in lawns in mixtures with Kentucky blue grass



POA CANBYI (Scribn.) Piper

Canby blue grass



Growth habit	erect, tufted, perennial, with fibrous roots
Blade	to 4 mm wide, 4 to 20 cm long, flat to folded, linear, boat-shaped at tip; scabrous on upper surface with scarious margins; smooth and prominently veined below; green and glaucous; folded at emergence
Sheath	compressed, keeled above, split, scabrous, light or yellowish green at base; margins hyaline
Ligule	to 5 mm long, membranous, acute, entire
Collar	divided or continuous, pale or yellowish green, often indistinct
Auricles	absent
Where found	throughout area in moist, slightly to moderately saline meadows



POA COMPRESSA L.

Canada blue grass



- Growth habit erect or rarely decumbent, sod-forming, perennial, with rhizomes
- Blade to 5 mm wide, 2 to 15 cm long, flat to folded, acuminate, tip boat-shaped, keeled below, broadest at base, bluish green, smooth or scabrous; veins not very prominent; margins scarious and scabrous; folded at emergence
- Sheath strongly compressed and sharply keeled, split, glabrous, green- or purplish-tinged at base; margins hyaline
- Ligule to 2 mm long, membranous, truncate or emarginate, entire, greenish
- Collar divided, narrow, glabrous, light or yellowish green
- Auricles absent
- Where found throughout area in moist meadows on poor soils; relatively abundant on sandy soils in eastern Saskatchewan and western Manitoba
- Remarks stem very flat, cannot be rolled between thumb and fingers



POA CUSICKII Vasey

early blue grass, Cusick blue grass


Growth habit	erect, tufted, perennial, with fibrous roots
Blade	to 3 mm wide, 5 to 15 cm long, usually folded and bristle-like, with boat-shaped tip, keeled, scabrous on margins, scabrous or smooth on surfaces; ridged on outer surface, with two prominent white veins resembling midrib, and continuing down sheath; folded at emergence
Sheath	compressed, sharply keeled, split, ridged, scabrous or smooth; margins overlapping, hyaline
Ligule	to 1 mm long, membranous, truncate, entire or slightly lacerate, very short
Collar	indistinct, very narrow, pale green
Auricles	absent
Where found	throughout area on moderately dry prairie



POA GLAUCIFOLIA Scribn. & Will.

glaucous blue grass



- Growth habit erect, loosely tufted, perennial, with few slender rhizomes
- Blade to 4 mm wide, 5 to 20 cm long, flat to folded, with boat-shaped tip, glabrous and glaucous on both surfaces, distinctly veined and prominently midribbed on lower surface; margins very scarious and scaberulous; folded at emergence
- Sheath slightly compressed, split, distinctly veined and somewhat keeled above, glabrous, often purplish at base; margins hyaline
- Ligule to 3 mm long, membranous, acute, entire
- Collar continuous or divided, light or yellowish green, often indistinct in younger shoots
- Auricles absent
- Where found throughout area in moist places but not abundant



POA PALUSTRIS L.

fowl blue grass



- Growth habit tall, loosely tufted, perennial, with fibrous roots
Blade to 4 mm wide, 7 to 15 cm long, lax, flat or V-shaped, with boat-shaped tip, broad at base; acuminate, scabrous on scarious margins and on both surfaces; mid-vein and two lateral veins evident on lower surface; folded at emergence
Sheath flattened and keeled, split, distinctly veined, smooth or slightly scabrous, pale green or purplish-tinged at decumbent base; margins overlapping, hyaline
Ligule to 4 mm long, membranous, acute, entire or lacerate
Collar distinct, divided, pale green, glabrous
Auricles absent
Where found throughout area in marshy places



POA PRATENSIS L.

Kentucky blue grass


- Growth habit erect, perennial, with long creeping rhizomes forming a dense sod
- Blade to 5 mm wide, 5 to 40 cm long, linear with boat-shaped tip, flat to folded, keeled, distinctly veined, smooth or scabrous on both surfaces, lower one usually smooth; margins slightly scabrous; dark green; folded at emergence
- Sheath compressed but not sharply keeled, split, distinctly veined, glabrous, dark green; margins hyaline
- Ligule to 1 mm long, membranous, truncate, entire
- Collar narrow, continuous, yellowish green, sometimes indistinct
- Auricles absent
- Where found throughout area on moist prairie in fertile soils; seeded in lawns and golf courses; becomes abundant in overgrazed fescue prairie
- Remarks naturally adventive from Europe about 1700; can be distinguished from *P. compressa* by its deeper green foliage, longer and parallel-sided blades, shorter ligule, and rounder stem



POA SANDBERGII Vasey

Sandberg's blue grass




Growth habit	low, erect, densely tufted, perennial, with fibrous roots
Blade	to 2 mm wide, 4 to 12 cm long, flat to folded, twisted, with boat-shaped tips, prominently veined, scabrous on both surfaces and on margins; folded at emergence
Sheath	compressed, split, smooth or slightly scabrous, prominently veined, pale or purplish at base; margins overlapping, hyaline
Ligule	to 3 mm long, membranous, acute, entire or lacerate
Collar	indistinct, divided, narrow, light green
Auricles	absent
Where found	throughout area on dry prairie
Remarks	a grass that matures its seed before July



POA TRIVIALIS L.

rough blue grass, rough-stalked meadow grass



- Growth habit perennial, tufted, with stolons and fibrous roots
Blade to 4.5 mm wide, 7 to 15 cm long, flat, acuminate, tip narrowly boat-shaped, slightly retrorsely scabrous on upper surface, glossy and keeled on lower surface; margins scabrous; bright green; folded at emergence
Sheath compressed and sharply keeled, generally scabrous, green or purple-tinted, split only partway
Ligule to 3 mm long, membranous, acute, entire or ciliate
Collar broad, distinct, glabrous, divided by midrib
Auricles absent
Where found occasionally seeded in moist to wet pastures in eastern regions
Remarks this grass resembles other species of *Poa* but can be distinguished by its scabrous sheath and the glossy undersurface of the blade



PUCCINELLIA NUTTALLIANA (Schultes) Hitchc.

Nuttall's alkali grass



Growth habit	erect to spreading, tufted, perennial, with fibrous roots
Blade	to 3 mm wide, 5 to 18 cm long, flat or involute, glaucous, very scabrous above, smooth below; margins slightly scabrous; veins indistinct; rolled at emergence
Sheath	round, split, smooth; margins overlapping, scarious
Ligule	to 5 mm long, membranous, acute to obtuse, continuous with margins of sheath
Collar	to 1 mm wide, not well defined
Auricles	absent
Where found	throughout area on dry to wet saline or alkaline soils
Remarks	often growing with <i>Distichlis stricta</i>



SCHEDONNARDUS PANICULATUS (Nutt.) Trel.

tumble grass

- Growth habit low, tufted, perennial, with fibrous roots
Blade to 2 mm wide, 2 to 5 cm long, usually folded, twisted and wavy, scabrous above, smooth below, scabrous on scarious margins and midrib; light bluish green; folded at emergence
Sheath compressed, sharply keeled, split, distinctly veined, very short, usually glabrous; margins broad, hyaline
Ligule to 3 mm long, membranous, acute, very conspicuous, and continuous with sheath margins
Collar divided, narrow, whitish
Auricles absent
Where found southern part of area on open prairie, especially on abandoned land, and on sandy or other infertile soil





K. F. B. 1921

SCHIZACHNE PURPURASCENS (Torr.) Swallen

purple oat grass


- Growth habit perennial, with rhizomes, growing erect from a loosely tufted, decumbent base
- Blade to 6 mm wide, 2 to 20 cm long, flat to folded, acuminate, narrowed at base, distinctly veined, scaberulous below, smooth or scaberulous on upper surface and margins; rolled at emergence
- Sheath round or slightly flattened, closed in early stages but splitting with maturity, purplish at base, smooth or scaberulous
- Ligule to 1 mm long, membranous, truncate, lacerate
- Collar indistinct, narrow, continuous, light green
- Auricles absent
- Where found throughout area in damp woods



SCOLOCHLOA FESTUCACEA (Willd.) Link

spangletop, thatch grass, scotch grass



Growth habit	tall, erect, perennial, with stout rhizomes
Blade	to 10 mm wide, 10 to 35 cm long, flat to convolute, ridged and slightly scabrous above, glabrous and faintly nerved below, acuminate, light to yellowish green; margins narrowly scarious and scabrous; rolled at emergence
Sheath	round, split, prominently veined, glabrous, yellowish white when mature; margins overlapping, hyaline
Ligule	to 8 mm long, membranous, truncate, lacerate
Collar	divided or continuous, glabrous, yellowish or brownish green
Auricles	absent
Where found	throughout area in shallow fresh to brackish sloughs and along streams



SETARIA GLAUCA (L.) Beauv.

yellow foxtail



- Growth habit erect or decumbent, annual, with fibrous roots
- Blade to 12 mm wide, 5 to 25 cm long, flat or V-shaped, acuminate, twisted, soft, drooping, glaucous, green, distinctly veined, midribbed below; slightly scabrous or smooth on both surfaces, with long twisted hairs near base on upper surface; margins smooth or scabrous; rolled at emergence from a compressed shoot
- Sheath compressed, sharply keeled, split, distinctly veined, glabrous, pale green, sometimes tinged with purple at base; margins overlapping, hyaline
- Ligule to 1 mm long, a fringe of hairs fused at the base
- Collar continuous, smooth, yellowish green, reddish or purple
- Auricles absent
- Where found in waste places, gardens, and cultivated areas; not common
- Remarks a weedy species introduced from Europe



SETARIA VIRIDIS (L.) Beauv.

green foxtail



- Growth habit coarse, semierect, annual, branching at base, with fibrous roots
- Blade to 12 mm wide, 5 to 20 cm long, flat, acuminate, light green, nodding, distinctly but finely veined, with prominent midvein below; margins serrulate-scabrous; rolled at emergence from a slightly compressed shoot
- Sheath slightly compressed, split, light green or purplish at base, glabrous or appressed pubescent; margins overlapping, inner margin hyaline, outer margin ciliate
- Ligule to 2 mm long, a fringe of hairs fused at base, with longer hairs at edges of collar margins
- Collar continuous, yellowish green or red-tinged, pubescent along margins
- Auricles absent
- Where found common in waste places, gardens, and cultivated fields
- Remarks a weedy species introduced from Europe



SITANION HYSTRIX (Nutt.) J. G. Smith

squirreltail


- Growth habit erect, rather short, densely tufted, perennial, with fibrous roots
- Blade to 3 mm wide, 2 to 20 cm long, flat to involute, acuminate, distinctly veined and finely pubescent on both surfaces; midvein prominent below; narrowly scarious; margins scabrous; rolled at emergence
- Sheath round, split partway down, somewhat keeled at upper end, indistinctly veined, glabrous or softly puberulent; margins hyaline
- Ligule to 1 mm long, membranous, truncate, lacerate, sometimes with wavy margins
- Collar continuous, light green, glabrous, sometimes indistinct
- Auricles clawlike when present
- Where found in very dry areas along the international boundary; rare
- Remarks may be mistaken for species of *Elymus* and *Hordeum*



SORGHASTRUM NUTANS (L.) Nash

Indian grass



- Growth habit erect, perennial, with scaly rhizomes
Blade to 10 mm wide, 10 to 50 cm long, flat, narrowed at base, acuminate, dull green to glaucous; veins and mid-rib prominent; margins and both surfaces scabrous; rolled at emergence
Sheath round or compressed near top, split, distinctly veined; upper sheaths usually glabrous, lower pubescent; often purplish-tinged below; margins hyaline
Ligule to 5 mm long, membranous, obtuse to truncate, lacerate, ciliate; margins often pubescent and veined, appearing to be projections of sheath margins
Collar continuous, glabrous, yellowish
Auricles absent
Where found eastern part of area on moist prairie; rare



SPARTINA GRACILIS Trin.

alkali cord grass



- Growth habit tall, erect, stiff-leaved, perennial, with scaly rhizomes
- Blade to 5 mm wide, 5 to 40 cm long, flat to convolute, stiff, coriaceous, acuminate, not prominently veined; upper surface and margins very scabrous, lower surface smooth; rolled at emergence
- Sheath round, split, distinctly veined, glabrous, yellowish green or purplish at base; margins overlapping, scarious
- Ligule to 2 mm long, a dense fringe of fine hairs
- Collar divided or continuous, broad, yellowish or brownish green, glabrous
- Auricles absent
- Where found throughout area in moist, moderately saline meadows and wet places
- Remarks this grass is more common in the western part of the area; *S. pectinata* is more common eastward



SPARTINA PECTINATA Link

prairie cord grass



Growth habit	tall, stout, perennial, with scaly rhizomes
Blade	to 15 mm wide, 10 to 60 cm long, flat to involute, coriaceous, long acuminate, distinctly veined above and below, very scabrous on margins and upper surface, smooth and shiny underneath, medium to light green, drooping; rolled at emergence
Sheath	round, split, distinctly veined, glabrous, pinkish at base; margins scarious
Ligule	to 4 mm long, a fringe of hairs, fused at the base
Collar	broad, continuous, glabrous, yellow or brownish green
Auricles	absent
Where found	eastern part of area, on moist prairie and in swamps; often abundant locally



SPHENOPHOLIS OBTUSATA (Michx.) Scribn.


prairie wedge grass



Growth habit	erect, tufted, perennial, with fibrous roots
Blade	to 5 mm wide, 3 to 20 cm long, flat, scabrous on both surfaces but more so on upper surface, acuminate, thin; margins very scabrous; midveins very fine below; rolled at emergence
Sheath	round, split, distinctly veined, glabrous or slightly scabrous; margins narrow, hyaline
Ligule	to 4 mm long, membranous, truncate, lacerate
Collar	narrow, continuous, or almost divided, often oblique, light or yellowish green
Auricles	absent
Where found	throughout area on moist and wet prairie; not common



SPOROBOLUS CRYPTANDRUS (Torr.) A. Gray

sand dropseed


- Growth habit erect or decumbent, tufted, perennial, with fibrous roots
- Blade  to 5 mm wide, 7 to 20 cm long, flat to involute or convolute, acuminate, distinctly veined; scabrous above and on scarious margins, smooth or scaberulous below, light green, fairly stiff; rolled at emergence
- Sheath round, split, prominently veined, glabrous, often purple-tinged at base; margins hyaline, ciliate
- Ligule to 1 mm long, a dense fringe of very fine silky hairs
- Collar broad, continuous, seldom divided, pubescent, yellow to dark green
- Auricles absent
- Where found throughout area on sandy soils
- Remarks often associated with *Calamovilfa longifolia* and *Oryzopsis hymenoides* in the sand dunes



SPOROBOLUS HETEROLEPIS (A. Gray) A. Gray

prairie dropseed


Growth habit	erect, densely tufted, perennial, occasionally with short rhizomes
Blade	to 3 mm wide, 5 to 45 cm long, flat to involute, acuminate, distinctly veined, scabrous on margins and upper surface, smooth and distinctly midribbed below, light green, erect or slightly drooping; rolled at emergence from a flattened shoot
Sheath	flattened, split, distinctly veined, glabrous or pubescent, white or purplish at base; margins hyaline
Ligule	to 0.5 mm long, membranous, truncate, ciliate
Collar	continuous, medium broad, pubescent on margins, yellowish green
Auricles	absent
Where found	eastern part of area on moist prairie



STIPA COMATA Trin. & Rupr.

needle-and-thread, common spear grass


Growth habit	erect, densely tufted, perennial, with fibrous roots
Blade	to 3 mm wide, 5 to 30 cm long, flat to involute or convolute, younger blades bristle-like, acuminate; ridged and scabrous on upper surface and margins, distinctly veined and smooth or scaberulous on lower surface; coriaceous, light green; rolled at emergence
Sheath	round or slightly compressed, split, prominently veined, glabrous; margins hyaline
Ligule	to 4 mm long, very conspicuous, membranous, obtuse or truncate, often split or frayed, continuous with sheath margins, sometimes ciliate
Collar	narrow, continuous, glabrous, light green, often oblique, usually indistinct in younger leaves
Auricles	absent
Where found	throughout area; the most common species on dry prairie



STIPA RICHARDSONII Link

Richardson's needle grass


- Growth habit erect, tufted, perennial, with fibrous roots
Blade to 3 mm wide, 5 to 25 cm long, involute, filiform, linear, acuminate; usually scabrous on inner surface and on narrowly scarious margins, smooth or scaberulous on outer surface; light green; veins indistinct on outer surface but prominent on inner; rolled at emergence
Sheath slightly flattened, split, distinctly veined, smooth or scabrous; margins hyaline
Ligule to 0.5 mm long, membranous, truncate
Collar indistinct, yellowish, continuous if evident
Auricles absent
Where found rare in Cypress Hills and Parklands; fairly common throughout the foothills of the Rocky Mountains
Remarks often mistaken for *S. curtisetia* (Hitc.) Barkworth, which has an erose ligule



STIPA SPARTEA Trin.

porcupine grass


- Growth habit erect, tufted, robust, perennial, with fibrous roots
- Blade to 5 mm wide, 5 to 40 cm long, flat to convolute; acuminate; ridged and scabrous on upper surface, distinctly veined and smooth on lower surface; midrib indistinct; coriaceous, light green, shiny, drooping; margins scarious; rolled at emergence
- Sheath round, split, prominently veined, white or purplish at base; margins hyaline, the outer usually ciliate
- Ligule to 5 mm long, membranous, obtuse or truncate, often split or irregular, finely ciliate
- Collar continuous, yellowish green, glabrous, often indistinct
- Auricles absent
- Where found eastern part of area on dry or moist prairie, as far west as the Qu'Appelle Valley
- Remarks *S. curtiseta* (Hitchc.) Barkworth has much the same growth habit but is less robust; the ligule is about one-half as long, the notch in the center forming two lateral lobes; the variety is found throughout the area



STIPA VIRIDULA Trin.

green needle grass



Growth habit	erect, tufted, perennial, with fibrous roots
Blade	to 5 mm wide, 10 to 25 cm long, flat to involute or convolute; prominently veined and scabrous on upper surface and to a lesser extent on lower surface; midrib prominent below; margins scabrous; rolled at emergence
Sheath	round, prominently veined, glabrous, split, villous near the throat; margins overlapping, scarious
Ligule	to 2 mm long, membranous, obtuse or truncate, entire
Collar	narrow, continuous, yellowish green, hairy all around but more so at margins
Auricles	absent
Where found	throughout area on moderately dry to moist, deep, fertile soils
Remarks	a variety from a plant improvement study has been named "green stipa grass," <i>Stipa viridula</i> Trin. cultivar



TRisetum SPICATUM (L.) Richt.

spike trisetum

- Growth habit erect, densely tufted, perennial, with fibrous roots
- Blade to 5 mm wide, 3 to 15 cm long, flat to involute, acuminate, distinctly veined; scaberulous and puberulent above and to a lesser extent below, rarely smooth; midrib prominent below; margins scabrous and scarious; rolled at emergence
- Sheath round, split only partway down, keeled at upper end, indistinctly veined, glabrous or puberulent; margins hyaline
- Ligule to 2 mm long, membranous, obtuse, lacerate
- Collar narrow, continuous, light green, often indistinct, with few hairs at margins
- Auricles absent
- Where found foothills of the Rocky Mountains, and northern parts of area in moist meadows



INDEX OF COMMON NAMES OF GRASSES
IN THE SECOND SECTION

Common name	Botanical name	Page
barley		
foxtail	<i>Hordeum jubatum</i> L.	148
wild	<i>Hordeum jubatum</i> L.	148
bent, creeping	<i>Agrostis stolonifera</i> L. var. <i>major</i> Farw.	52
bluejoint	<i>Agropyron smithii</i> Rydb.	40
bluestem		
big	<i>Andropogon gerardii</i> Vitman	58
little	<i>Andropogon scoparius</i> Michx.	60
brome		
California	<i>Bromus marginatus</i> Nees	74
downy	<i>Bromus tectorum</i> L.	78
fringed	<i>Bromus ciliatus</i> L.	70
northern awnless	<i>Bromus pumpellianus</i> Scribn.	76
smooth	<i>Bromus inermis</i> Leyss.	72
chess, downy	<i>Bromus tectorum</i> L.	78
cock's-foot	<i>Dactylis glomerata</i> L.	96
darnel, Persian	<i>Lolium persicum</i> Boiss. & Hohen.	154
dropseed		
prairie	<i>Sporobolus heterolepis</i> (A. Gray) A. Gray	230
sand	<i>Sporobolus cryptandrus</i> (Torr.) A. Gray	228
fescue		
bluebunch	<i>Festuca idahoensis</i> Elmer	132
meadow	<i>Festuca elatior</i> L. var. <i>pratensis</i> A. Gray	128
plains rough	<i>Festuca hallii</i> (Vasey) Piper	130
red	<i>Festuca rubra</i> L.	138
rough	<i>Festuca campestris</i> Rydb.	126
sheep	<i>Festuca ovina</i> L.	136
six-weeks	<i>Festuca octoflora</i> Walt.	134
foxtail		
green	<i>Setaria viridis</i> (L.) Beauv.	216
meadow	<i>Alopecurus pratensis</i> L.	56
short-awned	<i>Alopecurus aequalis</i> Sobol.	54
yellow	<i>Setaria glauca</i> (L.) Beauv.	214
grama		
blue	<i>Bouteloua gracilis</i> (H.B.K.) Lag. ex Steud.	68
side-oat	<i>Bouteloua curtipendula</i> (Michx.) Torr.	66
grass		
alkali	<i>Distichlis stricta</i> (Torr.) Rydb.	108
alkali cord	<i>Spartina gracilis</i> Trin.	222
annual blue	<i>Poa annua</i> L.	188
annual meadow	<i>Poa annua</i> L.	188
awned wheat	<i>Agropyron subsecundum</i> (Link) Hitchc.	44
barnyard	<i>Echinochloa crusgalli</i> (L.) Beauv.	110
bluebunch wheat	<i>Agropyron spicatum</i> (Pursh) Scribn. & Smith	42
brook	<i>Catabrosa aquatica</i> (L.) Beauv.	92
Canada blue	<i>Poa compressa</i> L.	192

Common name	Botanical name	Page
grass		
Canadian rice	<i>Oryzopsis canadensis</i> (Poir.) Torr.	168
canary	<i>Phalaris canariensis</i> L.	180
Canby blue	<i>Poa canbyi</i> (Scribn.) Piper	190
cane	<i>Phragmites communis</i> Trin.	186
common spear	<i>Stipa comata</i> Trin. & Rupr.	232
couch	<i>Agropyron repens</i> (L.) Beauv.	38
crested wheat	<i>Agropyron cristatum</i> (L.) Gaertn.	30
Cusick blue	<i>Poa cusickii</i> Vasey	194
desert salt	<i>Distichlis stricta</i> (Torr.) Rydb.	108
dwarf oat	<i>Danthonia unispicata</i> (Thurb.) Munro ex Macoun	104
early blue	<i>Poa cusickii</i> Vasey	194
false buffalo	<i>Munroa squarrosa</i> (Nutt.) Torr.	164
flyaway	<i>Agrostis scabra</i> Willd.	48
fowl blue	<i>Poa palustris</i> L.	198
fowl manna	<i>Glyceria striata</i> (Lam.) Hitchc.	142
glaucous blue	<i>Poa glaucifolia</i> Scribn. & Will.	196
green needle	<i>Stipa viridula</i> Trin.	238
green stipa	<i>Stipa viridula</i> Trin. cultivar	238
hair	<i>Agrostis scabra</i> Willd.	48
Hooker's oat	<i>Helictotrichon hookeri</i> (Scribn.) Henr.	144
Indian	<i>Sorghastrum nutans</i> (L.) Nash	220
Indian rice	<i>Oryzopsis hymenoides</i> (Roem. & Schult.) Ricker	170
intermediate wheat	<i>Agropyron intermedium</i> (Host) Beauv.	36
June	<i>Koeleria gracilis</i> Pers.	150
Kentucky blue	<i>Poa pratensis</i> L.	200
low spear	<i>Poa annua</i> L.	188
lyme	<i>Elymus virginicus</i> L.	124
marsh reed	<i>Calamagrostis canadensis</i> (Michx.) Beauv.	80
narrow reed	<i>Calamagrostis neglecta</i> (Ehrh.) Gaertn. Mey. & Schreb.	86
northern reed	<i>Calamagrostis inexpansa</i> A. Gray	82
northern rice	<i>Oryzopsis pungens</i> (Torr.) Hitchc.	172
northern wheat	<i>Agropyron dasystachyum</i> (Hook.) Scribn.	32
Nuttall's alkali	<i>Puccinellia nuttalliana</i> (Schultes) Hitchc.	206
one-spike oat	<i>Danthonia unispicata</i> (Thurb.) Munro ex Macoun	104
orchard	<i>Dactylis glomerata</i> L.	96
Parry oat	<i>Danthonia parryi</i> Scribn.	100
perennial rye	<i>Lolium perenne</i> L.	152
pine	<i>Calamagrostis rubescens</i> Buckl.	88
plains reed	<i>Calamagrostis montanensis</i> Scribn.	84
porcupine	<i>Stipa spartea</i> Trin.	236
poverty oat	<i>Danthonia spicata</i> (L.) Beauv. ex Roem. & Schult.	102
prairie cord	<i>Spartina pectinata</i> Link	224
prairie wedge	<i>Sphenopholis obtusata</i> (Michx.) Scribn.	226
purple oat	<i>Schizachne purpurascens</i> (Torr.) Swallen	210
quack	<i>Agropyron repens</i> (L.) Beauv.	38
reed canary	<i>Phalaris arundinacea</i> L.	178
Richardson's		
needle	<i>Stipa richardsonii</i> Link	234
rough blue	<i>Poa trivialis</i> L.	204
rough hair	<i>Agrostis scabra</i> Willd.	48

Common name	Botanical name	Page
grass		
rough-stalked		
meadow	<i>Poa trivialis</i> L.	204
salt	<i>Distichlis stricta</i> (Torr.) Rydb.	108
Sandberg's blue	<i>Poa sandbergii</i> Vasey	202
sand reed	<i>Calamovilfa longifolia</i> (Hook.) Scribn.	90
scotch	<i>Scolochloa festucacea</i> (Willd.) Link	212
scratch	<i>Muhlenbergia asperifolia</i> (Nees & Mey.) Parodi	156
slender wheat	<i>Agropyron trachycaulum</i> (Link) Malte	46
slender wood	<i>Cinna latifolia</i> (Trev.) Griseb.	94
slough	<i>Beckmannia syzigachne</i> (Steud.) Fern.	64
sweet	<i>Hierochloa odorata</i> (L.) Beauv.	146
switch	<i>Panicum virgatum</i> L.	176
tall manna	<i>Glyceria grandis</i> S. Wats.	140
tall wheat	<i>Agropyron elongatum</i> (Host) Beauv.	34
thatch	<i>Scolochloa festucacea</i> (Willd.) Link	212
tickle	<i>Agrostis scabra</i> Willd.	48
timber oat	<i>Danthonia intermedia</i> Vasey	98
tufted hair	<i>Deschampsia caespitosa</i> (L.) Beauv.	106
tumble	<i>Schedonnardus paniculatus</i> (Nutt.) Trel.	208
western rye	<i>Agropyron trachycaulum</i> (Link) Malte	46
western wheat	<i>Agropyron smithii</i> Rydb.	40
white-grained		
mountain rice	<i>Oryzopsis asperifolia</i> Michx.	166
wild oat	<i>Danthonia intermedia</i> Vasey	98
witch	<i>Panicum capillare</i> L.	174
muhly		
marsh	<i>Muhlenbergia racemosa</i> (Michx.) B.S.P.	160
mat	<i>Muhlenbergia richardsonis</i> (Trin.) Rydb.	162
prairie	<i>Muhlenbergia cuspidata</i> (Torr.) Rydb.	158
needle-and-thread	<i>Stipa comata</i> Trin. & Rupr.	232
redtop	<i>Agrostis stolonifera</i> L.	50
reed, common	<i>Phragmites communis</i> Trin.	186
rye		
Altai wild	<i>Elymus angustus</i> Trin.	112
blue wild	<i>Elymus glaucus</i> Buckl.	118
Canada wild	<i>Elymus canadensis</i> L.	114
giant wild	<i>Elymus cinereus</i> Scribn. & Merr.	116
hairy wild	<i>Elymus innovatus</i> Beal	120
Macoun's wild	<i>Agrohordeum macounii</i> (Vasey) Lepage	28
nodding wild	<i>Elymus canadensis</i> L.	114
Russian wild	<i>Elymus junceus</i> Fisch.	122
tufted wild	<i>Agrohordeum macounii</i> (Vasey) Lepage	28
Virginia wild	<i>Elymus virginicus</i> L.	124
spangletop	<i>Scolochloa festucacea</i> (Willd.) Link	212
squirreltail	<i>Sitanion hystrix</i> (Nutt.) J. G. Smith	218
three-awn, red	<i>Aristida longiseta</i> Steud.	62
timothy	<i>Phleum pratense</i> L.	184
alpine	<i>Phleum alpinum</i> L.	182
trisetum, spike	<i>Trisetum spicatum</i> (L.) Richt.	240

